


Lake Education and Planning Services, LLC
221B 2nd Street
Chetek, Wisconsin 54728

ROUND TRADE LAKE, BURNETT COUNTY

2018-2022 APMP
WDNR WBIC: 2640100

Prepared by: Dave Blumer, Lake Educator
November 22, 2017



RTLIA
Frederic, WI 54837

Distribution List

No. of Copies	Sent to
2	Vicki Breault, Vice-President Round-Trade Lake Improvement Association ?? ??
1	Pamela Toshner Wisconsin Department of Natural Resources 810 W. Maple Street Spooner, WI 54801

Table of Contents

INTRODUCTION	11
ROUND-TRADE LAKE IMPROVEMENT ASSOCIATION	13
PUBLIC PARTICIPATION AND STAKEHOLDER INPUT	14
OVERALL MANAGEMENT GOAL	15
EURASIAN WATERMILFOIL (EWM)	15
CURLY-LEAF PONDWEED (CLP)	16
ADDITIONAL MANAGEMENT GOALS FROM THE 2011 APM PLAN	19
WISCONSIN’S AQUATIC PLANT MANAGEMENT STRATEGY	20
LAKE INVENTORY	22
PHYSICAL CHARACTERISTICS	22
CRITICAL HABITAT	23
WATER QUALITY	23
FISHERIES AND WILDLIFE	26
ATTRIBUTES TO HELP MAINTAIN A HEALTHY LAKE AND WATERSHED	27
WETLANDS	27
SOILS	28
COARSE WOODY HABITAT (WOLTER, 2012)	29
SHORELANDS	30
<i>Protecting Water Quality</i>	31
<i>Natural Shorelands Role in Preventing AIS</i>	31
<i>Threats to Shorelands</i>	31
<i>Shoreland Preservation and Restoration</i>	32
AIS MANAGEMENT, 2007-2017	33
2010 AND 2016 WHOLE LAKE POINT INTERCEPT AQUATIC PLANT SURVEYS	35
<i>Simpson’s Diversity Index:</i>	36
<i>Floristic Quality Index (FQI)</i>	36
WILD RICE	38
AQUATIC INVASIVE SPECIES (AIS)	39
NON-NATIVE, AQUATIC INVASIVE PLANT SPECIES	39
<i>EWM</i>	39
<i>Curly-leaf Pondweed</i>	40
<i>Purple Loosestrife</i>	41
<i>Reed Canary Grass</i>	42
NON-NATIVE AQUATIC INVASIVE ANIMAL SPECIES	44
<i>Chinese Mystery Snails</i>	44
<i>Rusty Crayfish</i>	45
<i>Zebra Mussels</i>	45
AIS PREVENTION STRATEGY	47
MANAGEMENT ALTERNATIVES	48
NO MANAGEMENT	48
HAND-PULLING/MANUAL REMOVAL	49
DIVER ASSISTED SUCTION HARVESTING	50
MECHANICAL REMOVAL	51

<i>Large-scale Mechanical Harvesting</i>	51
<i>Small-Scale Cutting with removal</i>	52
BOTTOM BARRIERS AND SHADING	53
DREDGING	53
DRAWDOWN	53
BIOLOGICAL CONTROL	54
<i>EWM Weevils</i>	54
<i>Galerucella beetles</i>	55
<i>Other Biological Controls</i>	55
<i>Native Plant Restoration</i>	56
CHEMICAL CONTROL	56
<i>How Chemical Control Works</i>	57
<i>Efficacy of Aquatic Herbicides</i>	58
<i>Micro and Small-scale Herbicide Application</i>	58
<i>Large-scale Herbicide Application</i>	59
<i>Whole-Lake, And/Or Epilimnion Application</i>	59
<i>Effects of Whole-Lake Treatments on Native Aquatic Plant Species</i>	60
<i>Pre and Post Treatment Aquatic Plant Surveying</i>	60
<i>Chemical Concentration Testing</i>	61
HERBICIDE USE IN ROUND LAKE	61
MANAGEMENT DISCUSSION	63
EWM	63
CLP	63
<i>Application of Aquatic Herbicides</i>	63
AQUATIC PLANT SURVEYING	64
OTHER AIS MONITORING AND MANAGEMENT	64
COARSE WOODY HABITAT	64
SHORELAND IMPROVEMENT	65
WATER QUALITY	65
2018-2022 AQUATIC PLANT MANAGEMENT GOALS, OBJECTIVES, AND ACTIONS	66
IMPLEMENTATION AND EVALUATION	69
WISCONSIN DEPARTMENT OF NATURAL RESOURCES GRANT PROGRAMS	70
AIS PREVENTION AND CONTROL GRANTS	70
<i>Education, Prevention and Planning Projects</i>	70
<i>Established Population Control Projects</i>	71
<i>Maintenance and Containment Projects</i>	71
LAKE MANAGEMENT PLANNING GRANTS	71
<i>Small Scale Lake Management Projects</i>	72
<i>Large Scale Lake Management Projects</i>	72
LAKE PROTECTION GRANTS	73
<i>Healthy Lakes Projects</i>	73

Figures

Figure 1 – Trade River Watershed (SC10)	11
Figure 2 – Trade River Map – Round Lake (circled in red)	12
Figure 3 – 2010 (Polk County) Summer EWM; 2016 (ERS) Summer EWM	15
Figure 4 - 2012 and 2016 CLP Point-Intercept Survey Results	16
Figure 5 - 2012 and 2016 CLP Point-Intercept Survey Results	17
Figure 6 - 2016 CLP Point-Intercept Survey and Bed-mapping	18
Figure 7 – Rake Fullness Ratings	18
Figure 8 - Watershed Land Use for Round Lake, Burnett County (LEAPS, 2016)	22
Figure 9 – Trophic status in lakes	24
Figure 10 - Secchi Depth, Total Phosphorus, and Chlorophyll Graphs	25
Figure 11 - Round Lake Wetlands (Wisc. Wetlands Inventory December 30, 2016)	28
Figure 12 - Hydrologic Soil Group Classification in the Round Lake Watershed	29
Figure 13 - Coarse woody habitat-Fishsticks projects	30
Figure 14 - Healthy, AIS Resistant Shoreland (left) vs. Shoreland in Poor Condition	32
Figure 15 - 2012 fall EWM beds (left) and 2017 fall EWM beds (right)	34
Figure 16 – Changes in aquatic plant species identified during the 2010 and 2016 Point-intercept Aquatic Plant Surveys	36
Figure 17 - Wild Rice on Clam Lake in Burnett County (Photos by John Haack)	38
Figure 18 - EWM	40
Figure 19 - CLP Plants and Turions	41
Figure 20 - Purple Loosestrife	42
Figure 21 - Reed Canary Grass	43
Figure 22 - Chinese Mystery Snails	44
Figure 23 - Rusty Crayfish and identifying characteristics	45
Figure 24 - Zebra Mussels	47
Figure 25 - Aquatic vegetation manual removal zone	49
Figure 26 - DASH - Diver Assisted Suction Harvest (Aquacleaner Environmental, http://www.aquacleaner.com/index.html); Many Waters, LLC)	50
Figure 27 - EWM Weevil (https://klsa.wordpress.com/published-material/milfoil-weevil-guide/)	54
Figure 28 - Galerucella beetle	55
Figure 29 - Lake-wide (whole-lake) dissipation of aquatic herbicides in Mixed and Stratified Lakes (Carlson, 2015). Inset: Summer thermal stratification.	60

Tables

Table 1 – 2011-2017 EWM Management on Round Lake	17
Table 2 - Physical Characteristics of Round Lake in Burnett County.....	23
Table 3 - Management on Round Lake, 2007-2017	33
Table 4 – Herbicide treatments on Round Lake from 2012-2017	33
Table 5 - 2010 (Polk County) and 2016 (ERS) Point-intercept Aquatic Plant Survey Statistics	35
Table 6 – Measurements of Aquatic Plant Community Health	37
Table 7 - EWM Distribution based on Fall Bed-mapping Surveys.....	63

Appendices

Appendix A – NR 109	
Appendix B – NR 107	
Appendix C – WDNR Dredging Checklist	
Appendix D – Shoreland and Shallows Habitat Monitoring Field Protocol	
Appendix E – Round Lake Goals, Objectives, and Actions	
Appendix F – Funding and Implementation Matrix	
Appendix G – Calendar of Actions	

AQUATIC PLANT MANAGEMENT PLAN-ROUND LAKE

PREPARED FOR THE ROUND-TRADE LAKE IMPROVEMENT ASSOCIATION

INTRODUCTION

Round Lake is a 208 acre lake in Trade Lake Township in Burnett County. Round Lake is located in the Trade River watershed (Figure 1). The Trade River Watershed is approximately 124,754 acres in size and contains 167 miles of streams and rivers, 2,902 acres of lakes and 21,757 acres of wetlands. The watershed is dominated by forest (46%), grassland (19%) and wetlands (17%), and is ranked medium for nonpoint source issues affecting streams.




Figure 1 – Trade River Watershed (SC10)

Within the watershed, the Trade River begins in Polk County near Luck, WI, flows north in Burnett County and loops back to the south into the northwest of Polk County and then discharges into the St. Croix River (Figure 2). The Trade River flows through a chain of four lakes: Long Trade (Polk County), Round, Little Trade, and Big Trade (Burnett County). Round Lake is the second lake in a chain that the Trade River flows through on its way to the St. Croix River.

These four lakes have been united under a common lake association for many years. The Round - Trade Lake Improvement Association (RTLIA) has been an active and enthusiastic leader in the pioneering efforts at lake management in Wisconsin. The four lakes along the Trade River have exhibited signs of excess fertility for decades. The signs of eutrophication are evident on these waters but a definitive nutrient and hydraulic budget has not been documented. A feasibility study to evaluate the hydraulic and nutrient loading to the lakes as well as in-lake monitoring to determine recycling and profile characteristics is a high priority for these lakes.




Figure 2 – Trade River Map – Round Lake (circled in red)

Round Lake is considered a “Deep Lowland” lake under the state's Natural Community Determinations. Deep lowland lakes are generally deep enough to stratify during the summer season, have water draining from them, and may have water entering them from upstream. Stated lake uses for Round Lake are fishing and swimming. These uses are not being met, and the lake was added to the Wisconsin Impaired Waters list in 2012 for Total Phosphorus, Excess Algal Growth, and Eutrophication. Round Lake was first listed as an impaired water in 1998 for mercury build up in fish tissue caught in the lake. During the assessment process for the 2018 impaired waters list, water quality data continued to exceed thresholds for recreation use and fish and aquatic life.

WDNR Watersheds <http://dnr.wi.gov/water/waterDetail.aspx?key=16674>

Eurasian watermilfoil (EWM) was identified in Round Lake in 2003. Curly-leaf pondweed (CLP), another aquatic invasive species (AIS) was officially recognized in 2008, but has likely been in the lake much longer than that.

ROUND-TRADE LAKE IMPROVEMENT ASSOCIATION

The Lake Improvement Association is a non-profit, state incorporated, association comprised of Round Lake, Big and Little Trade Lakes, Long Trade Lake, and Spirit Lake (Spirit Lake is no longer part of the Association). These lakes are located in both Burnett and Polk counties in Wisconsin. The official name is the Round-Trade Lake Improvement Association, Inc., however, it is referred to as the Lake Improvement Association, or more recently the RTLIA. The RTLIA was originally incorporated March 26, 1974, by residents of Round and Trade Lakes with its official offices located in the Township of Trade Lake. The RTLIA is comprised of officers: President, Vice-President, Secretary-Treasurer. Each of the officers is elected through the nomination and vote process of the association's members. There are four board members representing each of the lakes in the association and there is also a lake chairman from each of the four lakes. Regular funding is generated from annual membership dues, with additional funding coming from grants obtained through the state of Wisconsin. Four annual association open meetings are held where members of the association or the public can present ideas or concerns.

The following is a list of actions the RTLIA have implemented over the years. This list should not be considered a complete list, but does reflect the reasons for the efforts made.

- Installed larger culverts on Trade Lake in order to control the lake water levels and reduce flooding of the lakeshore.
- Initiated a plan to control the number of rough fish in Round and Trade Lakes.
- Funded aquatic plant management actions in an effort to control the weed growth for recreational users of the lakes.
- Worked with the DNR to mark hazardous areas with buoys on Trade and Round Lakes.
- Successfully defeated the development of an asphalt plant that was to be located on the Trade River, just north of Round Lake.
- Fought to deter the development of back lot access to Round Lake.
- Introduced weevils into Long Trade Lake in an effort to control the spread of Eurasian watermilfoil.
- Completed Aquatic Plant Management Plans for all four lakes
- Applied for and received State grant funding for planning and control of EWM and CLP
- Completed a study of the water chemistry, macrophytes, and lake sediment of Round Lake.
- Is working to complete a study of the watershed of Long Trade Lake
- Participated in annual state conservation conferences, contributed financially to local causes, and disseminated information to local lakeshore owners on various conservation methods.

Membership in the RTLIA requires payment of dues in the sum of \$30.00 annually. Completed membership forms and investment can be given to any of the officers, board members, lake chairman, or it can be mailed. A current listing of all the officers, board members, and chairman is available by visiting the RTLIA web site at www.tradelakeassoc.org. Minutes from meetings, additional information about the RTLIA, and many other resources are available on the webpage. The RTLIA is also on Facebook. Some current and future goals for the RTLIA include but are not limited to the following:

- Complete water quality studies of the lakes in the Trade River system
- Develop long-term strategies to maintain and improve the water quality of the lakes in the Trade River system
- Control or reduce the spread of exotic plant species in the lakes and on the surrounding lakeshores.
- Increase active membership in the Lake Improvement Association.

PUBLIC PARTICIPATION AND STAKEHOLDER INPUT

OVERALL MANAGEMENT GOAL

The overall management goal for Round Lake is to maintain EWM at low levels that do not interfere with lake use. For the purposes of this management plan, that level is less than three acres as identified annually during a late summer littoral (plant growing) zone point-intercept (PI) survey. A secondary goal is to reduce the amount of dense growth CLP in the lake by 50% in five years as measured by spring CLP bed-mapping and reduce the number of turions in the sediment by 50% after three years of active management.

In the previous Trade Lakes Aquatic Plant Management Plan (2011), management objectives for EWM in the entire system included:

1. 75% reduction in dense growth EWM over five years;
2. Preventing EWM from spreading into Big Trade Lake;
3. Preventing EWM from leaving the lake via boat traffic; and
4. Getting property owners to take action to control EWM near their docks and shoreland.

Of these objectives, only the first and last can be said to have been fully reached. Based on 2016 totals, EWM in all four lakes combined showed an 80% reduction. However, it did spread from Little Trade Lake into Big Trade Lake during this time period. Property owners on Round Lake have taken an active interest in controlling EWM by their own accord. Public access on Round Lake is available on the southwest side of the lake, with 120 hours of watercraft inspection recorded in the WDNR SWIMS database since 2015.

EURASIAN WATERMILFOIL (EWM)

Two whole lake summer surveys of the vegetation in Round Lake have been completed. A summer point-intercept survey was completed in 2010 by Polk County, and a summer point-intercept survey was completed again in 2016 by Endangered Resource Sciences (ERS). During the 2010 survey, 15.5 acres of EWM was identified (Figure 3), and in the 2016 summer survey EWM covered approximately 7.7 acres.




Figure 3 – 2010 (Polk County) Summer EWM; 2016 (ERS) Summer EWM

CURLY-LEAF PONDWEED (CLP)

One of the goals in the 2011 Trade Lakes APM Plan was to document the extent of CLP in each of the four lakes, and then to create an addendum to the existing plan focused on control of CLP. This goal was met, with a CLP management plan for all four lakes being completed in 2012. Round Lake had the least amount of CLP when compared to the other three lakes in the system. During the 2012 cold-water/early season survey, CLP was identified at 9 points (Figure 4). During the 2016 cold-water/early season survey, CLP was identified at 21 points (Figure 4). CLP bed-mapping in 2012 only mapped 5 acres. In 2016 the area decreased to 1.2 acres, but this was after management using diquat, a contact herbicide that was applied to control EWM (Figure 4). During both surveys, CLP was present in the rest of the lake, but not common. CLP in the rest of the system does exceptionally well over thick organic muck. This type of bottom substrate is limited in Round Lake which could explain why there is not more CLP in the lake.

Once the extent of CLP in Round Lake was known, several scenarios for management were proposed. The expected scenario, which was first implemented in 2013, focused on treating all of the EWM in the lake, but none of the CLP. Unlike the other three lakes, CLP was not targeted at all in the course of managing EWM in Round Lake. There may have been some incidental take on CLP (like in 2016), but it was not targeted directly.




Figure 4 - 2012 and 2016 CLP Point-Intercept Survey Results




Figure 5 - 2012 and 2016 CLP Point-Intercept Survey Results

EWM has been chemically treated every year since 2012 with mixed results (Table 1). In most years some combination of granular and liquid 2,4-D was used. In 2016, diquat, a faster acting contact herbicide, was used as it was thought that in past years water movement through the lake was diluting the herbicide applied making it less effective. The 2016 treatment was a complete waste of resources, as the diquat had no impact on the EWM in the lake. This was likely due to it being applied in deeper water than is customary for this herbicide. As a result the concentration reached was nowhere near enough to kill or even slow down the growth of EWM (Table 1).

Table 1 – 2011-2017 EWM Management on Round Lake

AIS Management on Round Lake, 2011-2017							
Task	2011	2012	2013	2014	2015	2016	2017
APM Plan	X						X
AIS Control Grant		X	X	X	X	X	X
AIS Rapid Response Grant							
Spring EWM Treatment (acres)		5.96	15.74	5.68	4.08	4.13	4.84
Fall EWM Bed Mapping	X	X	X	X	X	X	X
EWM Physical Removal		X	X	X	X	X	X
CLP Bed Mapping		X				X	
Spring CLP Treatment (acres)							
Pre-treatment Plant Survey		X	X	X	X		
Post Treatment Plant Survey		X	X	X	X		
Whole-lake PI Survey		X				X	
X= Completed, P=Proposed							

CLP bed-mapping in 2016 only identified 1.2 acres of CLP left in the lake after treatment. All of it was considered moderate or dense with a rake fullness rating of 2 or 3 on a 1-3 scale (Figures 6 & 7). CLP management in 2018 and beyond will target both the CLP located in the same areas as proposed EWM treatments, and additional acreage along developed shores with the intent of reducing the total presence of CLP (actual plant acreage and turions) to 50% or less of 2016 levels within five years. A continued goal would be to reduce the total presence of CLP by at least another 50% within ten years. Turion reductions would be based off of 2018 survey numbers.




Figure 6 - 2016 CLP Point-Intercept Survey and Bed-mapping




<u>Rating</u>	<u>Coverage</u>	<u>Description</u>
1		A few plants on rake head
2		Rake head is about 1/2 full Can easily see top of rake head
3		Overflowing Cannot see top of rake head

Figure 7 – Rake Fullness Ratings

ADDITIONAL MANAGEMENT GOALS FROM THE 2011 APM PLAN

Additional management recommendations made in the 2011 APM Plan included preventing the spread and introduction of other aquatic invasive species; preserving, protecting, and enhancing the lake's native aquatic plant community; and minimizing runoff of pollutants, nutrients, and sediment from the Trade River Watershed. In an effort to reach these goals and meet the objectives stated for them, the RTLIA continued participation in a purple loosestrife biological control program by raising beetles; sponsored several AIS educational events; participated in AIS monitoring; and encouraged property owners to leave native aquatic plants in place along their shorelines. Developing a nutrient budget and watershed management plan was also included as a management objective, however the process to begin doing this has only been started since 2016, and then only for Long Trade Lake at the top of the Trade River system that includes Long Trade, Round, Little Trade, and Big Trade Lakes.

WISCONSIN'S AQUATIC PLANT MANAGEMENT STRATEGY

The waters of Wisconsin belong to all people. Their management becomes a balancing act between the rights and demands of the public and those who own property on the water's edge. This legal tradition called the Public Trust Doctrine dates back hundreds of years in North America and thousands of years in Europe. Its basic philosophy with respect to the ownership of waters was adopted by the American colonies. The US Supreme Court has found that the people of each state hold the right to all their navigable waters for their common use, such as fishing, hunting, boating and the enjoyment of natural scenic beauty.

The Public Trust Doctrine is the driving force behind all management in Wisconsin lakes. Protecting and maintaining that resource for all of Wisconsin's people are at the top of the list in determining what is done and where. In addition to the Public Trust Doctrine, two other forces have converged that reflect Wisconsin's changing attitudes toward aquatic plants. One is a growing realization of the importance of a strong, diverse community of aquatic plants in a healthy lake ecosystem. The other is a growing concern over the spread of AIS, such as EWM. These two forces have been behind more recent changes in Wisconsin's aquatic plant management laws and the evolution of stronger support for the control of invasive plants.

To some, these two issues may seem in opposition, but on closer examination they actually strengthen the case for developing an APMPs as part of a total lake management picture. Planning is a lot of work, but a sound plan can have long-term benefits for a lake and the community living on and using the lake.

The impacts of humans on Wisconsin's waters over the past five decades have caused public resource professionals in Wisconsin to evolve a certain philosophy toward aquatic plant management. This philosophy stems from the recognition that aquatic plants have value in the ecosystem, as well as from the awareness that, sometimes, excessive growth of aquatic plants can lessen our recreational opportunities and our aesthetic enjoyment of lakes. In balancing these, sometimes competing objectives, the Public Trust Doctrine requires that the State's public resource professionals be responsible for the management of fish and wildlife resources and their sustainable use to benefit all Wisconsin citizens. Aquatic plants are recognized as a natural resource to protect, manage, and use wisely.

Aquatic plant protection begins with human beings. We need to work to maintain good water quality and healthy native aquatic plant communities. The first step is to limit the amount of nutrients and sediment that enter the lake. There are other important ways to safeguard a lake's native aquatic plant community. They may include developing motor boat ordinances that prevent the destruction of native plant beds and reduce shoreline erosion and sediment disturbance caused by boat wakes, limiting aquatic plant removal activities, designating certain plant beds as critical habitat sites and preventing the spread of non-native, invasive plants, such as EWM.

If plant management is needed, it is usually in lakes that humans have significantly altered. If we discover how to live on lakes in harmony with natural environments and how to use aquatic plant management techniques that blend with natural processes rather than resist them, the forecast for healthy lake ecosystems looks bright. To assure no harm is done to the lake ecology, it is important that plant management is undertaken as part of a long range and holistic plan.

In many cases, the development of long-term, integrated aquatic plant management strategies to identify important plant communities and manage nuisance aquatic plants in lakes, ponds or rivers is required by the State of Wisconsin. To promote the long-term sustainability of our lakes, the State of Wisconsin endorses the development of APMPs and supports that work through various grant programs.

There are many techniques for the management of aquatic plants in Wisconsin. Often management may mean protecting desirable aquatic plants by selectively hand pulling the undesirable ones. Sometimes more

intensive management may be needed such as using harvesting equipment, herbicides or biological control agents. These methods require permits and extensive planning. Often using an Integrated Pest Management (IPM) strategy that incorporates multiple management actions/alternatives works the best.

While limited management on individual properties is generally permitted, it is widely accepted that a lake will be much better off if plants are considered on a whole lake scale. This is routinely accomplished by lake organizations or units of government charged with the stewardship of individual lakes.

LAKE INVENTORY

In order to make recommendations for aquatic plant and lake management, basic information about the water body of concern is necessary. A basic understanding of physical characteristics including size and depth, critical habitat, water quality, water level, fisheries and wildlife, wetlands and soils is needed to make appropriate recommendations for improvement.

PHYSICAL CHARACTERISTICS

Round Lake (WBIC 2640100) is a hard water drainage lake located in south central Burnett County. It is the second in a chain of four lakes along the Trade River. According to the Wisconsin Department of Natural Resources (WDNR), Round Lake has a surface area of 208 acres and a maximum depth of 27 feet.

The primary land use in the Round Lake watershed is forest land which occupies about 42% of the 35,595 acre watershed with agricultural uses (i.e. row crops, pasture, etc.) at around 38%. Development accounts for about 8% of the land cover with the heaviest development (land that is covered by greater than 50% impervious surfaces) contributing less than 1%. Of this developed area, very little is found around Round Lake. Wetlands and undeveloped grasslands account for just under 9% of the land use in the Round Lake watershed while open water covers the remaining 3% (Figure 8, Table 2).




Figure 8 - Watershed Land Use for Round Lake, Burnett County (LEAPS, 2016)

Table 2 - Physical Characteristics of Round Lake in Burnett County

Lake Characteristics	
Lake Area (acres)	207.3
Watershed Area (acres)	35,596.60
Maximum Depth (feet)	27
Mean Depth (feet)	15
Miles of Shoreline	5
Lake Type	Drainage

Watershed Land Use	
Open Water	1,230.70
Forest	15,074.90
Agriculture (Crops and Pasture)	13,574.80
Grassland/ Scrub	994.7
Heavily Developed	207
Lightly Developed	2,467.40
Wetlands	2,045.10
Total	35,596.60

Land cover and land use management practices have a strong influence on water quality. Increases in impervious surfaces, such as roads, rooftops and compacted soils, associated with residential and agricultural land uses can reduce or prevent the infiltration of runoff. This can lead to an increase in the amount of rainfall runoff that flows directly into Round Lake as well as into the Trade River. The removal of riparian, i.e., near shore, vegetation causes an increase in the amount of nutrient-rich soil particles transported directly to the lake during rain events.

Round Lake has one active resort on the lake.

CRITICAL HABITAT

Every body of water has areas of aquatic vegetation that offers critical or unique fish and wildlife habitat. Such areas can be identified by the WDNR and identified as Sensitive Areas per Ch. NR 107. Although Round Lake has not been surveyed by the WDNR for sensitive areas, there are areas of the lake that should be left in an undisturbed state to provide aquatic habitat and ecosystem services necessary for a healthy lake. Aquatic habitat areas provide the basic needs (e.g. habitat, food, nesting areas) for waterfowl, fish, and wildlife. Disturbance to these areas during mechanical harvesting should be avoided or minimized and chemical treatment is generally not allowed. Areas of rock and cobble substrate with little or no fine sediment are considered high quality walleye spawning habitat. No dredging, structures, or deposits should occur in these sensitive areas.

WATER QUALITY

One of the most commonly used metrics of water quality is the trophic state of a lake. The trophic state is defined as the total load of biomass in a waterbody at any given time (Carlson & Simpson, 1996). To determine the trophic state of any given lake, the Tropic State Index (TSI) is generally used. This index uses the three main variables of Secchi depth, total phosphorus, and chlorophyll concentration. TSI values are technically limitless, but when applied, they almost always fall between 0 and 100. To make sense of these values, they are broken into different trophic states. The four main trophic states are oligotrophic, mesotrophic, eutrophic, and hypereutrophic (Figure 9). Oligotrophic lakes are usually very clear, clean lakes with low nutrient levels. Mesotrophic lakes are moderately clear with some nutrients and more plants present within the system. Eutrophic lakes have excess nutrients that support a great deal of algae growth, and may have a large aquatic plant community. Hypereutrophic lakes are very green with dense algae and limited plant growth. Round Lake is considered to be a eutrophic lake.




Figure 9 – Trophic status in lakes

Round Lake has been classified as an “Impaired Water” by the WDNR. The Clean Water Act mandates that every even numbered year the WDNR (and all other states) must prepare a list of waters that do not meet the water quality standards established by the state. A waterbody is defined as impaired if it is shown that one or more pollutant criteria are met and if it “does not support full use by humans, wildlife, fish, and other aquatic life” (WDNRa, 2015). Round Lake has received this designation due to excess algal growth, total phosphorus content, and eutrophication. All water quality data in this section were retrieved from the WDNR Round Lake webpage.

Water quality data has been collected on Round Lake since 1986 by Citizen Lake Monitoring Network (CLMN) volunteers. Citizen lake monitoring plays a critical role in collecting data to determine water quality trends over time. All of the data collected on Round Lake is collected from the deep hole site in the center of the lake. Average water clarity, or Secchi depth, data range from 1.9 feet in 1986 up to 13 feet in 1999. The average annual Secchi depth for Round Lake since data has been collected is 5 feet (Figure 7).

Average summer total phosphorus data range from 44-µg/l in 2013 to 141-µg/l in 2012. The overall average total phosphorus from 2005-2016 is 71-µg/l, which suggests Round Lake is eutrophic based on phosphorus values (Figure 10). The average total phosphorus in 2012 was almost twice than is normal for Round Lake. However there was only one measurement taken in 2012, and field notes taken by the CLMN volunteers suggest that Round Lake experienced a very large rain a few days before this sample was taken. This rain event could have resulted in higher levels of runoff entering the lake causing a spike in phosphorus levels. Even when this abnormally high measurement is removed from the data, the yearly phosphorus still averages 64-µg/l making Round Lake a eutrophic lake.

Average summer chlorophyll concentration data ranges from 25- $\mu\text{g}/\text{l}$ in 2013 to 59.85- $\mu\text{g}/\text{l}$ in 2005. The average chlorophyll concentration for Round Lake was 43- $\mu\text{g}/\text{l}$ which classifies Round Lake as eutrophic. The unusually low chlorophyll concentrations in 2013 could be explained by the unusual conditions described in the field notes. These notes described 2013 as an uncharacteristically cold year with the ice remaining in place longer than usually and summer temperatures being below average.

This data suggests that the water quality in Round Lake has not gotten any worse, and may actually have gotten better since the last APM plan was completed in 2011.




Figure 10 - Secchi Depth, Total Phosphorus, and Chlorophyll Graphs

FISHERIES AND WILDLIFE

Until recent years, Round Lake had been stocked fairly regularly with walleye since 1973. The most recent stocking consisted of 7,493 walleye fingerlings (4-6 inch) in 2011. Aside from walleye, panfish, largemouth bass, and northern pike are common while smallmouth bass are present.

According to the Natural Heritage Inventory (NHI), there are two protected species within the same township as Round Lake. The brittle prickly-pear (*Opuntia fragilis*) is a terrestrial plant that is found in dry sandy areas. While this threatened plant is not found in close proximity to Round Lake, it should be considered for any projects that concern the Round Lake watershed. The second protected species found near Round Lake is Blanding's turtle (*Emydoidea blandingii*) which is not listed as threatened or endangered, but is considered a species of special concern, and is listed as a protected wild animal under NR10.02. It is fairly likely that Blanding's turtles are present within and around Round Lake. Minimizing impact to these turtles should be considered when planning management.

WETLANDS

A wetland is an area where water is at, near or above the land surface long enough to be capable of supporting aquatic or hydrophytic vegetation and which has soils indicative of wet conditions. Wetlands have many functions which benefit the ecosystem surrounding Round Lake. Wetlands with a higher floral diversity of native species support a greater variety of native plants and are more likely to support regionally scarce plants and plant communities. Wetlands provide fish and wildlife habitat for feeding, breeding, resting, nesting, escape cover, travel corridors, spawning grounds for fish, and nurseries for mammals and waterfowl.

Wetlands also provide flood protection within the landscape. Due to the dense vegetation and location within the landscape, wetlands are important for retaining stormwater from rain and melting snow moving towards surface waters and retaining floodwater from rising streams. This flood protection minimizes impacts to downstream areas. Wetlands provide water quality protection because wetland plants and soils have the capacity to store and filter pollutants ranging from pesticides to animal wastes.

Wetlands also provide shoreline protection to Round Lake because shoreline wetlands act as buffers between land and water. They protect against erosion by absorbing the force of waves and currents and by anchoring sediments. This shoreline protection is important in waterways where boat traffic, water current, and wave action cause substantial damage to the shore. Wetlands also provide groundwater recharge and discharge by allowing the surface water to move into and out of the groundwater system. The filtering capacity of wetland plants and substrates help protect groundwater quality. Wetlands can also stabilize and maintain stream flows, especially during dry months. Aesthetics, recreation, education and science are also all services wetlands provide. Wetlands contain a unique combination of terrestrial and aquatic life and physical and chemical processes.

Round Lake only has one wetland that borders the lake. However this wetland is located along the southwestern shoreline where the Trade River enters Round Lake. Most of the water that enters Round Lake will pass through this wetland complex which can help reduce the nutrients entering the lake from the rest of the watershed. This wetland can also provide quality habitat for wildlife and help reduce the erosion caused by the river.




Figure 11 - Round Lake Wetlands (Wisc. Wetlands Inventory December 30, 2016)

SOILS

Soils are classified into hydrologic soil groups to indicate their potential for producing runoff. Group A soils have a high infiltration rate which makes the potential amount of runoff very low. These soils are, generally very sandy and allow water to pass through unimpeded. Conversely, group D soils have a very low infiltration rate making their runoff potential fairly high. Group D soils are generally very dense with high amounts of organic material. This causes water to move slowly through group D soils often resulting in standing water on flat surfaces and flowing water over sloped surfaces. Most of the soils within the Round Lake watershed fall into groups C and C/D (NRCSa, 2016). These soils are very good for agriculture, but they also have a fairly high runoff potential. If managed appropriately, the runoff can be minimized in these areas.




Figure 12 - Hydrologic Soil Group Classification in the Round Lake Watershed

COARSE WOODY HABITAT (WOLTER, 2012)

Coarse woody habitat (CWH) in lakes is classified as trees, limbs, branches, roots, and wood fragments at least 4 inches in diameter that enter a lake by natural (beaver activity, toppling from ice, wind, or wave scouring) or human means (logging, intentional habitat improvement, flooding following dam construction). CWH in the littoral or near-shore zone serves many functions within a lake ecosystem including erosion control, as a carbon source, and as a surface for algal growth which is an important food base for aquatic macro invertebrates. Presence of CWH has also been shown to prevent suspension of sediments, thereby improving water clarity. CWH serves as important refuge, foraging, and spawning habitat for fish, aquatic invertebrates, turtles, birds, and other animals. The amount of littoral CWH occurring naturally in lakes is related to characteristics of riparian forests and likelihood of toppling. However, humans have also had a large impact on amounts of littoral CWH present in lakes through time. During the 1800's the amount of CWH in northern lakes was increased beyond natural levels as a result of logging practices. But time changes in the logging industry and forest composition along with increasing shoreline development have led to reductions in CWH present in many northern Wisconsin lakes.

CWH is often removed by shoreline residents to improve aesthetics or select recreational opportunities (swimming and boating). Jennings et al. (2003) found a negative relationship between lakeshore development and the amount of CWH in northern Wisconsin lakes. Similarly, Christensen et al. (1996) found a negative correlation between density of cabins and CWH present in Wisconsin and Michigan lakes. While it is difficult to make precise determinations of natural densities of CWH in lakes it is believed that the value is likely on the scale of hundreds of logs per mile. The positive impact of CWH on fish communities have been well documented by researchers, making the loss of these habitats a critical concern. One study determined that black crappie selected nesting sites that were usually associated with woody debris, silty substrate, warmer water, and protected from wind and waves (Pope & Willis, 1997).

Fortunately, remediation of this habitat type is attainable on many waterbodies, particularly where private landowners and lake associations are willing to partner with county, state, and federal agencies. Large-scale CWH projects are currently being conducted by lake associations and local governments with assistance from the WDNR where hundreds of whole trees are added to the near-shore areas of lakes. For more information on this process visit: <http://dnr.wi.gov/topic/fishing/outreach/fishsticks.html> (last accessed on 12-29-2016).

Small-scale CWH projects, more commonly referred to as “fishsticks,” can also be done by individual property owners, and are eligible for grant assistance through the WDNR Healthy Lakes program. This program is intended to help individual property owners make a positive impact on their lake’s ecosystem through small-scale projects such as fishsticks (Figure 13).



Figure 13 - Coarse woody habitat-Fishsticks projects

SHORELANDS

How the shoreline of a lake is managed can have big impacts on the water quality and health of that lake. Natural shorelines prevent polluted runoff from entering lakes, help control flooding and erosion, provide fish and wildlife habitat, may make it harder for AIS to establish themselves, muffle noise from watercraft, and preserve privacy and natural scenic beauty. Many of the values lake front property owners appreciate and enjoy about their properties - natural scenic beauty, tranquility, privacy, relaxation - are enhanced and preserved with good shoreland management. And healthy lakes with good water quality translate into healthy lake front property values.

Shorelands may look peaceful, but they are actually the hotbed of activity on a lake. 90% of all living things found in lakes - from fish, to frogs, turtles, insects, birds, and other wildlife - are found along the shallow margins and shores. Many species rely on shorelands for all or part of their life cycles as a source for food, a place to sleep, cover from predators, and to raise their young. Shorelands and shallows are the spawning grounds for fish, nesting sites for birds, and where turtles lay their eggs. There can be as much as 500% more species diversity at the water's edge compared to adjoining uplands.

Lakes are buffered by shorelands that extend into and away from the lake. These shoreland buffers include shallow waters with submerged plants (like coontail and pondweeds), the water's edge where fallen trees and emergent plants like rushes might be found, and upward onto the land where different layers of plants (low ground cover, shrubs, trees) may lead to the lake. A lake's littoral zone is a term used to describe the shallow water area where aquatic plants can grow because sunlight can penetrate to the lake bottom. Shallow lakes might be composed entirely of a littoral zone. In deeper lakes, plants are limited where they can grow by how deeply light can penetrate the water.

Shorelands are critical to a lake’s health. Activities such as replacing natural vegetation with lawns, clearing brush and trees, importing sand to make artificial beaches, and installing structures such as piers, can cause water

quality decline and change what species can survive in the lake. In addition to being potentially damaging, some of these undertakings require permits and approval. Most changes to lakebed exposed by fluctuating water levels (removal of sediments, additions of beach sand, etc.), often require permits and approval. The only exceptions to this are manual removal of a 30 foot corridor of native plants or the removal of non-native invasive plants. These regulations have been put in place to encourage property owners to responsibly manage their shorelands to improve and maintain the quality of the lake as a whole.

PROTECTING WATER QUALITY

Shoreland buffers slow down rain and snow melt (runoff). Runoff can add nutrients, sediments, and other pollutants into lakes, causing water quality declines. Slowing down runoff will help water soak (infiltrate) into the ground. Water that soaks into the ground is less likely to damage lake quality and recharges groundwater that supplies water to many of Wisconsin's lakes. Slowing down runoff water also reduces flooding, and stabilizes stream flows and lake levels.

Shoreland wetlands act like natural sponges trapping nutrients where nutrient-rich wetland sediments and soils support insects, frogs, and other small animals eaten by fish and wildlife.

Shoreland forests act as filters, retainers, and suppliers of nutrients and organic material to lakes. The tree canopy, young trees, shrubs, and forest understory all intercept precipitation, slowing runoff, and contributing to water infiltration by keeping the soil's organic surface layer well-aerated and moist. Forests also slow down water flowing overland, often capturing its sediment load before it can enter a lake or stream. In watersheds with a significant proportion of forest cover, the erosive force of spring snow melts is reduced as snow in forests melts later than snow on open land, and melt water flowing into streams is more evenly distributed. Shoreland trees grow, mature, and eventually fall into lakes where they protect shorelines from erosion, and are an important source of nutrients, minerals and wildlife habitat.

NATURAL SHORELANDS ROLE IN PREVENTING AIS

In addition to removing essential habitat for fish and wildlife, clearing native plants from shorelines and shallow waters can open up opportunities for invasive species to take over. Like tilling a home garden to prepare it for seeding, clearing shoreland plants exposes bare earth and removes the existing competition (the cleared shoreland plants) from the area. Nature fills a vacuum. While the same native shoreland plants may recover and reclaim their old space, many invasive species possess "weedy" traits that enable them to quickly take advantage of new territory and out-compete natives.

The act of weeding creates continual disturbance, which in turn benefits plants that behave like weeds. The modern day practice of mowing lawns is an example of keeping an ecosystem in a constant state of disturbance to the benefit of invasive species like turf grass, dandelions, and clover, all native to Europe. Keeping shoreline intact is a good way to minimize disturbance and minimize opportunities for invasive species to gain a foothold.

THREATS TO SHORELANDS

When a landowner develops a waterfront lot, many changes may take place including the addition of driveways, houses, decks, garages, sheds, piers, rafts and other structures, wells, septic systems, lawns, sandy beaches and more. Many of these changes result in the compaction of soil and the removal of trees and native plants, as well as the addition of impervious (hard) surfaces, all of which alter the path that precipitation takes to the water.

Building too close to the water, removing shoreland plants, and covering too much of a lake shore lot with hard surfaces (such as roofs and driveways) can harm important habitat for fish and wildlife, send more nutrient and sediment runoff into the lake, and cause water quality decline.

Changing one waterfront lot in this fashion may not result in a measurable change in the quality of the lake or stream. But cumulative effects when several or many lots are developed in a similar way can be enormous. A lake's response to stress depends on what condition the system is in to begin with, but bit by bit, the cumulative effects of tens of thousands of waterfront property owners "cleaning up" their shorelines, are destroying the shorelands that protect their lakes. Increasing shoreline development and development throughout the lake's watershed can have undesired cumulative effects.

SHORELAND PRESERVATION AND RESTORATION

If a native buffer of shoreland plants exists on a given property, it can be preserved and care taken to minimize impacts when future lake property projects are contemplated. If a shoreline has been altered, it can be restored. Shoreline restoration involves recreating buffer zones of natural plants and trees. Not only do quality wild shorelines create higher property values, but they bring many other values too. Some of these are aesthetic in nature, while others are essential to a healthy ecosystem. Healthy shorelines mean healthy fish populations, varied plant life, and the existence of the insects, invertebrates and amphibians which feed fish, birds and other creatures. Figure 14 shows the difference between a natural and unnatural shoreline adjacent to a lake home. More information about healthy shorelines can be found at the following website: <http://wisconsinlakes.org/index.php/shorelands-a-shallows> (last accessed 12-27-2016).




Figure 14 - Healthy, AIS Resistant Shoreland (left) vs. Shoreland in Poor Condition

Some of the property surrounding Round Lake has been left natural. However, there are some areas that have been heavily developed. In these areas, the natural shoreland buffer has been disturbed which contributes negatively to lake health by preventing rainwater and snowmelt from being filtered by the native vegetation before entering the lake. Improvements to the shoreline in these areas can help lessen human impact on the lake, and many of these projects are fairly easy to do.

AIS MANAGEMENT, 2007-2017

Management in Round Lake has been focused on a well-established population of EWM. While the three other lakes on the Trade River also struggle with CLP, its impact on Round Lake has been minimal. CLP populations seem to be staying near the Trade River inlet on the southern shore which is allowing management efforts to be directed at EWM. While CLP is still being monitored, there has been no need to actively manage is. Table 3 breaks down yearly management efforts from 2007 to 2017.

Table 3 - Management on Round Lake, 2007-2017

AIS Management on Round Lake, 2007-2017											
Task	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
APM Plan					X						X
AIS Control Grant						X	X	X	X	X	X
AIS Rapid Response Grant											
Spring EWM Treatment (acres)						5.96	15.74	5.7	4.08	4.13	4.84
Fall EWM Bed Mapping					X	X	X	X	X	X	X
EWM Physical Removal						X	X	X	X	X	X
CLP Bed Mapping						X				X	
Spring CLP Treatment (acres)											
Pre-treatment Plant Survey						X	X	X	X	X	
Post Treatment Plant Survey						X	X	X	X	X	
Whole-lake PI Survey				X		X				X	
X= Completed, P=Proposed											

EWM was first found in Round Lake in 2003, and was left essentially unmanaged until the initial APM Plan was written in 2011 which paved the way for management to begin in 2012. This allowed EWM to gain a foothold in most of the littoral zone within Round Lake by the time the 2012 fall bed mapping surveys were done (Figure 15). Since treatment began in 2012, the EWM population within Round Lake seems to be shrinking with the large-scale 2013 treatment seeming to have the biggest impact. The EWM treatments and the impacts on both AIS and native species are shown in Table 4.

Table 4 – Herbicide treatments on Round Lake from 2012-2017

Year	# of Beds	Total Area Treated (acres)	Range of Bed Size	Herbicide	Concentration	Results- AIS	Results- Native Plants
2012	7	5.96	0.24-1.55	Granular 2,4-D	0.92 acres @4.0 ppm 5.04 acres @ 3.0ppm	EWM- Significant decrease	Coontail, Wild celery, Water star-grass, and Sago pondweed- Significant increases. All others- No changes.
2013	21	15.74	0.11-2.13	Liquid 2,4-D	15.74 acres @ 3.0ppm	EWM- Significant decrease	Common waterweed, Wild celery, White water lily, and Water star-grass- Significant increases. All others- No changes
2014	5	5.68	0.24-1.79	Liquid 2,4-D Granular 2,4- D	5.44 acres @ 3.0 ppm 0.24 acres @ 4.0 ppm	EWM- less than ideal due to high water	Wild celery, White water lily, Slender naiad- Significant increases, All other- No changes
2015	8	4.08	0.11-1.48	Liquid 2,4-D Granular 2,4- D	3.03 acres @ 3.5 ppm 1.05 acres @ 3.5-4.0 ppm	EWM- Significant decrease	Common waterweed, Claspig-leaf pondweed, and wild celery- Significant increases, All other- No changes
2016	4	4.16	0.15-2.9	Diquat	2 gal/acre*ft	No decrease in EWM	Significant decrease in CLP
2017	4	4.84	0.8-1.59	Liquid 2,4-D	1.39 acres @ 3.5ppm 3.45 acres @ 4.0ppm	EWM-Significant decrease	Results pending




Figure 15 - 2012 fall EWM beds (left) and 2017 fall EWM beds (right)

2010 AND 2016 WHOLE LAKE POINT INTERCEPT AQUATIC PLANT SURVEYS

A prerequisite to updating the APMP for Round Lake was to compare how the lake’s vegetation had changed since the last point intercept survey. In 2010, a warm-water, whole-lake, point-intercept survey of aquatic plants was completed by the Polk County Land and Water Conservation Department. This survey was repeated in 2016 by Endangered Resource Sciences (ERS) on July 12. During the 2010 survey EWM was documented at thirteen points with one other point having a visual record of the plant. In 2016, twenty-one points were identified with EWM during the summer survey, with an additional 16 points having a visual record. Part of the increase is due to the lack of EWM that was afforded by the use of diquat in the spring of the year. In the two surveys, the littoral (plant growing) zone was different with the 2010 littoral zone extending to about 15-ft. and the 2016 littoral zone only extended to about 9-ft. Aquatic plant data was only collected between 3 and 15 feet of water in the 2010 survey, so is less dependable than the data collected in 2016. Based on the 2016 survey, the littoral zone of Round Lake covers about 40 acres or about 38.5% of the total lake surface area.

Table 5 - 2010 (Polk County) and 2016 (ERS) Point-intercept Aquatic Plant Survey Statistics

SUMMARY STATS: 2010 & 2016 Summer PI Survey	2010	2016
Total number of sites visited	333	1000
Total number of sites with vegetation	112	127
Total number of sites shallower than maximum depth of plants	318	179
Frequency of occurrence at sites shallower than maximum depth of plants	35.22	70.95
Simpson Diversity Index	0.87	0.91
Maximum depth of plants (ft)**	15.50	8.50
Number of sites sampled using rake on Rope (R)	8	0
Number of sites sampled using rake on Pole (P)	326	288
Average number of all species per site (shallower than max depth)	0.78	2.08
Average number of all species per site (veg. sites only)	2.22	2.94
Average number of native species per site (shallower than max depth)	0.74	1.88
Average number of native species per site (veg. sites only)	2.17	2.74
Species Richness	19	26
Species Richness (including visuals)	22	28
Species Richness (including visuals and boat survey)	27	38
Mean depth of plants (ft)	5.55	2.79
Median depth of plants (ft)	4.00	2.00
Mean rake fullness (veg. sites only)	1.49	2.02




Figure 16 – Changes in aquatic plant species identified during the 2010 and 2016 Point-intercept Aquatic Plant Surveys

Three measurements of the health of the aquatic plant community outside of these survey statistics are the Simpson’s Diversity Index (SDI), Floristic Quality Index (FQI), and Coefficient of Conservatism.

SIMPSON’S DIVERSITY INDEX:

A diversity index allows the entire plant community at one location to be compared to the entire plant community at another location. It also allows the plant community at a single location to be compared over time thus allowing a measure of community degradation or restoration at that site. With Simpson’s Diversity Index, the index value represents the probability that two individual plants (randomly selected) will be different species. The index values range from 0 -1 where 0 indicates that all the plants sampled are the same species to 1 where none of the plants sampled are the same species. The greater the index value, the higher the diversity in a given location. Although many natural variables like lake size, depth, dissolved minerals, water clarity, mean temperature, etc. can affect diversity, in general, a more diverse lake indicates a healthier ecosystem. Perhaps most importantly, plant communities with high diversity also tend to be more resistant to invasion by exotic species.

The SDI in 2010 was 0.87. In 2016 the value increased to 0.91. These values are similar to Long Trade Lake, and slightly higher than Little Trade Lake, also in the Trade River system (Table 6).

FLORISTIC QUALITY INDEX (FQI)

This index measures the impact of human development on a lake’s aquatic plants. The 124 species in the index are assigned a Coefficient of Conservatism (C) which ranges from 1-10. The higher the value assigned, the more likely the plant is to be negatively impacted by human activities relating to water quality or habitat modifications. Plants with low values are tolerant of human habitat modifications, and they often exploit these changes to the point where they may crowd out other species. The FQI is calculated by averaging the

conservatism value for each native index species found in the lake during the point-intercept survey, and multiplying it by the square root of the total number of plant species (N) in the lake. Statistically speaking, the higher the index value, the healthier the lake's aquatic plant community is assumed to be. Nichols (1999) identified four eco-regions in Wisconsin: Northern Lakes and Forests, North Central Hardwood Forests, Driftless Area and Southeastern Wisconsin Till Plain. He recommended making comparisons of lakes within ecoregions to determine the target lake's relative diversity and health. Round Lake is in the North Central Hardwood Forests Ecoregion.

In 2010, 16 native index species were identified in the rake during the point-intercept survey. They produced a mean C of 5.1 and a FQI of 20.3. In 2016, 21 native index plants were identified in the rake during the point-intercept survey. They produced a mean C of 5.0 and a FQI of 22.9. Nichols (1999) reported an average mean C for the North Central Hardwood Forests Region of 5.6 putting Round Lake below average for this part of the state. The FQI was just slightly above the median FQI of 20.9 for the North Central Hardwood Forests (Nichols 1999) (Table 6).

Although several broadleaf pondweed species (flat-stem pondweed, sago pondweed, Fries pondweed, and water stargrass) appear to be down in 2016, this is likely due to the application of diquat in the late spring of the year to control EWM. Unfortunately, not much EWM was killed, but a limited amount of CLP and these other species were. It remains to be seen if the four native plant species will recover from the diquat treatment. It is expected that they will, given that diquat usually only kills what it comes in contact with. All of these plant species regrow from seeds or rhizomes in the sediment.

Otherwise, based on the measurements of the health of the aquatic plant community, Round Lake hasn't changed much from 2010. Those changes that are documented are likely the result of the increased number of different aquatic plant species that were identified in the 2016 survey. As previously stated, these species are likely the result of a more experienced aquatic plant surveyor than they are the result of new plants in the lake.

Eurasian watermilfoil is down in the system, but it does not appear that the overall population can be brought down below several acres, even with annual treatments. CLP is increasing in the lake, and for that reason, any EWM management action should include CLP management at the same time.

Table 6 – Measurements of Aquatic Plant Community Health

Parameter	2010	2016
SDI	0.87	0.91
FQI	20.3	22.9
Mean C	5.1	5
# of Species	16	21

WILD RICE

According to the Great Lakes Indian Fish and Wildlife Commission (GLIFWC), Long Trade, Round, Little Trade and Big Trade lakes are not wild rice waters. Additionally, wild rice was not found during the aquatic plant surveys of the lakes or during the Sensitive Areas survey. Although wild rice is not present in these lakes, it warrants attention due to its ecologic and cultural significance and its abundance in nearby lakes and streams (for example, the Grettum Flowage, Rice Lake, Spirit Lake, and the Clam Lakes). Any activity included in a comprehensive lake or aquatic plant management plan that could potentially impact the growth of wild rice in any body of water that has in the past, currently has, or potentially could have wild rice in the future requires consultation with the Tribal Nations. This consultation is completed by the Department of Natural Resources during their review of lake management documents. When present in a lake, wild rice is afforded numerous protections due to its ecological and cultural significance and management is therefore focused on harvest goals and protection rather than removal.

Wild rice is an annual aquatic grass that produces seed that is a nutritious source of food for wildlife and people (Figure 20). As a native food crop, it has a tremendous amount of cultural significance to the Wisconsin and Minnesota Native American Nations. Wild rice pulls large amounts of nutrients from the sediment in a single year and the stalks provide a place for filamentous algae and other small macrophytes to attach and grow. These small macrophytes pull phosphorous in its dissolved state directly from the water. Wild rice can benefit water quality, provide habitat for wildlife, and help minimize substrate re-suspension and shoreland erosion.

In Wisconsin, wild rice has historically ranged throughout the state. Declines in historic wild rice beds have occurred statewide due to many factors, including dams, pollution, large boat wakes, and invasive plant species. Renewed interest in the wild rice community has led to large-scale restoration efforts to reintroduce wild rice in Wisconsin's landscape. There is the potential for planting wild rice at shoreline restoration and rehabilitation sites in the Trade Lakes system however this should not be done without full constituent support as the presence of wild rice will limit certain aquatic plant management actions. Extensive information is available on wild rice from GLIFWC and the WDNR.



Figure 17 - Wild Rice on Clam Lake in Burnett County (Photos by John Haack)

AQUATIC INVASIVE SPECIES (AIS)

Past invasive species monitoring efforts have identified several different plant and animal non-native, invasive species in Round Lake. Most of these species are considered aquatic, although some are also considered shoreland or wetland type invasive species.

NON-NATIVE, AQUATIC INVASIVE PLANT SPECIES

Eurasian watermilfoil (EWM) and curly-leaf pondweed (CLP) are the most problematic invasive species found on Long Trade Lake. In addition purple loosestrife and reed canary grass have been identified along the shores of Long Trade Lake. Purple loosestrife and reed canary grass are shoreland or wetland plants not generally problematic within the lake, but can be very problematic on the shores and in the wetlands adjacent to the lake. More information is given for each non-native species in the following sections (WDNRb, 2016).

EWM

EWM is a submersed aquatic plant native to Europe, Asia, and northern Africa (Figure 18). It is the only non-native milfoil in Wisconsin. Like the native milfoils, the Eurasian variety has slender stems whorled by submersed feathery leaves and tiny flowers produced above the water surface. The flowers are located in the axils of the floral bracts, and are either four-petaled or without petals. The leaves are threadlike, typically uniform in diameter, and aggregated into a submersed terminal spike. The stem thickens below the inflorescence and doubles its width further down, often curving to lie parallel with the water surface. The fruits are four-jointed nut-like bodies. Without flowers or fruits, EWM is difficult to distinguish from Northern water milfoil. EWM has 9-21 pairs of leaflets per leaf, while Northern milfoil typically has 7-11 pairs of leaflets. Coontail is often mistaken for the milfoils, but does not have individual leaflets.

EWM grows best in fertile, fine-textured, inorganic sediments. In less productive lakes, it is restricted to areas of nutrient-rich sediments. It has a history of becoming dominant in eutrophic, nutrient-rich lakes, although this pattern is not universal. It is an opportunistic species that prefers highly disturbed lake beds, lakes receiving nitrogen and phosphorous-laden runoff, and heavily used lakes. Optimal growth occurs in alkaline systems with a high concentration of dissolved inorganic carbon. High water temperatures promote multiple periods of flowering and fragmentation.

Unlike many other plants, EWM does not rely on seed for reproduction. Its seeds germinate poorly under natural conditions. It reproduces by fragmentation, allowing it to disperse over long distances. The plant produces fragments after fruiting once or twice during the summer. These shoots may then be carried downstream by water currents or inadvertently picked up by boaters. EWM is readily dispersed by boats, motors, trailers, bilges, live wells, and bait buckets; and can stay alive for weeks if kept moist.

Once established in an aquatic community, milfoil reproduces from shoot fragments and stolons (runners that creep along the lake bed). As an opportunistic species, EWM is adapted for rapid growth early in spring. Stolons, lower stems, and roots persist over winter and store the carbohydrates that help milfoil claim the water column early in spring, photosynthesize, divide, and form a dense leaf canopy that shades out native aquatic plants. Its ability to spread rapidly by fragmentation and effectively block out sunlight needed for native plant growth often results in monotypic stands. Monotypic stands of EWM provide only a single habitat, and threaten the integrity of aquatic communities in a number of ways; for example, dense stands disrupt predator-prey relationships by fencing out larger fish, and reducing the number of nutrient-rich native plants available for waterfowl.

Dense stands of EWM also inhibit recreational uses like swimming, boating, and fishing. Some stands have been dense enough to obstruct industrial and power generation water intakes. The visual impact that greets

the lake user on milfoil-dominated lakes is the flat yellow-green of matted vegetation, often prompting the perception that the lake is "infested" or "dead". Cycling of nutrients from sediments to the water column by EWM may lead to deteriorating water quality and algae blooms in infested lakes.



Figure 18 - EWM

Round Lake has a fully established population of EWM.

CURLY-LEAF PONDWEED

Round Lake has an established population of curly-leap pondweed. While CLP is not currently being aggressively managed, preventing its spread should still be considered when planning management actions for Round Lake.

Curly-leaf pondweed (CLP) is an invasive aquatic perennial that is native to Eurasia, Africa, and Australia (Figure 19). It was accidentally introduced to United States waters in the mid-1880s by hobbyists who used it as an aquarium plant. The leaves are reddish-green, oblong, and about 3 inches long, with distinct wavy edges that are finely toothed. The stem of the plant is flat, reddish-brown and grows from 1 to 3 feet long. CLP is commonly found in alkaline and high nutrient waters, preferring soft substrate and shallow water depths. It tolerates low light and low water temperatures. It has been reported in all of the Continental United States with Wisconsin having the highest density of CLP.

CLP spreads through burr-like winter buds (turions) (Figure 19), which are moved among waterways. These plants can also reproduce by seed, but this plays a relatively small role compared to the vegetative reproduction through turions. New plants form under the ice in winter, making curly-leaf pondweed one of the first nuisance aquatic plants to emerge in the spring. It becomes invasive in some areas because of its tolerance for low light and low water temperatures. These tolerances allow it to get a head start on and out-compete native plants in the spring. In mid-summer, when most aquatic plants are growing, CLP plants are dying off. Plant die-offs may result in a critical loss of dissolved oxygen. Furthermore, the decaying plants can increase nutrients which contribute to algal blooms, as well as create unpleasant stinking messes on beaches. CLP forms surface mats that interfere with aquatic recreation.



Figure 19 - CLP Plants and Turions

PURPLE LOOSESTRIFE

Purple loosestrife (Figure 20) is a perennial herb 3-7 feet tall with a dense bushy growth of 1-50 stems. The stems, which range from green to purple, die back each year. Showy flowers that vary from purple to magenta possess 5-6 petals aggregated into numerous long spikes, and bloom from August to September. Leaves are opposite, nearly linear, and attached to four-sided stems without stalks. It has a large, woody taproot with fibrous rhizomes that form a dense mat. By law, purple loosestrife is a nuisance species in Wisconsin. It is illegal to sell, distribute, or cultivate the plants or seeds, including any of its cultivars.

Purple loosestrife is a wetland herb that was introduced as a garden perennial from Europe during the 1800's. It is still promoted by some horticulturists for its beauty as a landscape plant, and by beekeepers for its nectar-producing capability. Currently, more than 20 states, including Wisconsin have laws prohibiting its importation or distribution because of its aggressively invasive characteristics. It has since extended its range to include most temperate parts of the United States and Canada. The plant's reproductive success across North America can be attributed to its wide tolerance of physical and chemical conditions characteristic of disturbed habitats, and its ability to reproduce prolifically by both seed dispersal and vegetative propagation. The absence of natural predators, like European species of herbivorous beetles that feed on the plant's roots and leaves, also contributes to its proliferation in North America.

Purple loosestrife was first detected in Wisconsin in the early 1930's, but remained uncommon until the 1970's. It is now widely dispersed in the state, and has been recorded in 70 of Wisconsin's 72 counties. Low densities in most areas of the state suggest that the plant is still in the pioneering stage of establishment. Areas of heaviest infestation are sections of the Wisconsin River, the extreme southeastern part of the state, and the Wolf and Fox River drainage systems.

This plant's optimal habitat includes marshes, stream margins, alluvial flood plains, sedge meadows, and wet prairies. It is tolerant of moist soil and shallow water sites such as pastures and meadows, although established plants can tolerate drier conditions. Purple loosestrife has also been planted in lawns and gardens, which is often how it has been introduced to many of our wetlands, lakes, and rivers.

Purple loosestrife can germinate successfully on substrates with a wide range of pH. Optimum substrates for growth are moist soils of neutral to slightly acidic pH, but it can exist in a wide range of soil types. Most seedling establishment occurs in late spring and early summer when temperatures are high.

Purple loosestrife spreads mainly by seed, but it can also spread vegetatively from root or stem segments. A single stalk can produce from 100,000 to 300,000 seeds per year. Seed survival is up to 60-70%, resulting in an extensive seed bank. Mature plants with up to 50 shoots grow over 2 meters high and produce more than two

million seeds a year. Germination is restricted to open, wet soils and requires high temperatures, but seeds remain viable in the soil for many years. Even seeds submerged in water can live for approximately 20 months. Most of the seeds fall near the parent plant, but water, animals, boats, and humans can transport the seeds long distances. Vegetative spread through local perturbation is also characteristic of loosestrife; clipped, trampled, or buried stems of established plants may produce shoots and roots. Plants may be quite large and several years old before they begin flowering. It is often very difficult to locate non-flowering plants, so monitoring for new invasions should be done at the beginning of the flowering period in mid-summer.

Any sunny or partly shaded wetland is susceptible to purple loosestrife invasion. Vegetative disturbances such as water drawdown or exposed soil accelerate the process by providing ideal conditions for seed germination. Invasion usually begins with a few pioneering plants that build up a large seed bank in the soil for several years. When the right disturbance occurs, loosestrife can spread rapidly, eventually taking over the entire wetland. The plant can also make morphological adjustments to accommodate changes in the immediate environment; for example, a decrease in light level will trigger a change in leaf morphology. The plant's ability to adjust to a wide range of environmental conditions gives it a competitive advantage; coupled with its reproductive strategy, purple loosestrife tends to create monotypic stands that reduce biotic diversity. Purple loosestrife displaces native wetland vegetation and degrades wildlife habitat. As native vegetation is displaced, rare plants are often the first species to disappear. Eventually, purple loosestrife can overrun wetlands thousands of acres in size, and almost entirely eliminate the open water habitat. The plant can also be detrimental to recreation by choking waterways.

Purple loosestrife has been identified in several locations around Round Lake.



Figure 20 - Purple Loosestrife

REED CANARY GRASS

Reed canary grass (Figure 21) is a large, coarse grass that reaches 2 to 9 feet in height. It has an erect, hairless stem with gradually tapering leaf blades 3 1/2 to 10 inches long and 1/4 to 3/4 inch in width. Blades are flat and have a rough texture on both surfaces. The lead ligule is membranous and long. The compact panicles are erect or slightly spreading (depending on the plant's reproductive stage), and range from 3 to 16 inches long with branches 2 to 12 inches in length. Single flowers occur in dense clusters in May to mid-June. They are green to purple at first and change to beige over time. This grass is one of the first to sprout in spring, and forms a thick rhizome system that dominates the subsurface soil. Seeds are shiny brown in color.

Both Eurasian and native ecotypes of reed canary grass are thought to exist in the U.S. The Eurasian variety is considered more aggressive, but no reliable method exists to tell the ecotypes apart. It is believed that the vast majority of our reed canary grass is derived from the Eurasian ecotype. Agricultural cultivars of the grass are widely planted.

Reed canary grass is a cool-season, sod-forming, perennial wetland grass native to temperate regions of Europe, Asia, and North America. The Eurasian ecotype has been selected for its vigor and has been planted throughout the U.S. since the 1800's for forage and erosion control. It has become naturalized in much of the northern half of the U.S., and is still being planted on steep slopes and banks of ponds and created wetlands.

Reed canary grass can grow on dry soils in upland habitats and in the partial shade of oak woodlands, but does best on fertile, moist organic soils in full sun. This species can invade most types of wetlands, including marshes, wet prairies, sedge meadows, fens, stream banks, and seasonally wet areas; it also grows in disturbed areas such as berms and spoil piles.

Reed canary grass reproduces by seed or creeping rhizomes. It spreads aggressively. The plant produces leaves and flower stalks for 5 to 7 weeks after germination in early spring and then spreads laterally. Growth peaks in mid-June and declines in mid-August. A second growth spurt occurs in the fall. The shoots collapse in mid to late summer, forming a dense, impenetrable mat of stems and leaves. The seeds ripen in late June and shatter when ripe. Seeds may be dispersed from one wetland to another by waterways, animals, humans, or machines.

This species prefers disturbed areas, but can easily move into native wetlands. Reed canary grass can invade a disturbed wetland in just a few years. Invasion is associated with disturbances including ditching of wetlands, stream channelization, and deforestation of swamp forests, sedimentation, and intentional planting. The difficulty of selective control makes reed canary grass invasion of particular concern. Over time, it forms large, monotypic stands that harbor few other plant species and are subsequently of little use to wildlife. Once established, reed canary grass dominates an area by building up a tremendous seed bank that can eventually erupt, germinate, and recolonize treated sites.

Reed canary grass can be found in the wetland that borders the southwestern shoreline of Round Lake. There are portions of this wetland that contain dense stands of reed canary grass, but most of the wetland has not become overrun with this grass.



Figure 21 - Reed Canary Grass

NON-NATIVE AQUATIC INVASIVE ANIMAL SPECIES

Several non-vegetative, aquatic, invasive species are in nearby lakes, but have not been identified in Round Lake. Two species, Chinese mystery snails and rusty crayfish have been seen on some parts of the Trade River. This means that these species could be in Round Lake, or they will have an easier time finding their way into Round Lake (WDNRb, 2016). It is important for lake property owners and users to be knowledgeable of these species in order to identify them if and when they show up in Round Lake.

CHINESE MYSTERY SNAILS

Chinese mystery snails and banded mystery snail have not been identified in Round Lake, but they have been spotted in the Trade River about 1.5 miles upstream of the Round Lake inlets.

The Chinese mystery snails and the banded mystery snails (Figure 22) are non-native snails that have been found in a number of Wisconsin lakes. There is not a lot yet known about these species, however, it appears that they have a negative effect on native snail populations. The mystery snail's large size and hard operculum (a trap door cover which protects the soft flesh inside), and their thick hard shell make them less edible by predators.

The female mystery snail gives birth to live crawling young. This may be an important factor in their spread as it only takes one impregnated snail to start a new population. Mystery snails thrive in silt and mud areas although they can be found in lesser numbers in areas with sand or rock substrates. They are found in lakes, ponds, irrigation ditches, and slower portions of streams and rivers. They are tolerant of pollution and often thrive in stagnant water areas. Mystery snails can be found in water depths of 0.5 to 5 meters (1.5 to 15 feet). They tend to reach their maximum population densities around 1-2 meters (3-6 feet) of water depth. Mystery snails do not eat plants. Instead, they feed on detritus and in lesser amounts on algae and phytoplankton. Thus removal of plants along the shoreline area will not reduce the abundance of mystery snails.

Lakes with high densities of mystery snails often see large die-offs of the snails. These die-offs are related to the lake's warming coupled with low oxygen (related to algal blooms). Mystery snails cannot tolerate low oxygen levels. High temperatures by themselves seem insufficient to kill the snails as the snails could move into deeper water.

Many lake residents are worried about mystery snails being carriers of the swimmer's itch parasite. In theory they are potential carriers, however, because they are an introduced species and did not evolve as part of the lake ecosystem, they are less likely to harbor the swimmer's itch parasites.



Figure 22 - Chinese Mystery Snails

RUSTY CRAYFISH

Rusty crayfish have not been identified within Round Lake, but there is an established population along most of the Trade River. Because of this, it is very likely that there are some rusty crayfish within Round Lake, but there is not an established population within the lake.

Rusty crayfish (Figure 23) live in lakes, ponds and streams, preferring areas with rocks, logs and other debris in water bodies with clay, silt, sand or rocky bottoms. They typically inhabit permanent pools and fast moving streams of fresh, nutrient-rich water. Adults reach a maximum length of 4 inches. Males are larger than females upon maturity and both sexes have larger, heartier, claws than most native crayfish. Dark “rusty” spots are usually apparent on either side of the carapace, but are not always present in all populations. Claws are generally smooth, with grayish-green to reddish-brown coloration. Adults are opportunistic feeders, feeding upon aquatic plants, benthic invertebrates, detritus, juvenile fish and fish eggs.

The native range of the rusty crayfish includes Ohio, Tennessee, Kentucky, Indiana, Illinois and the entire Ohio River basin. However, this species may now be found in Michigan, Massachusetts, Missouri, Iowa, Minnesota, New York, New Jersey, Pennsylvania, Wisconsin, New Mexico and the entire New England state area (except Rhode Island). The Rusty crayfish has been a reported invader since at least the 1930’s. Its further spread is of great concern since the prior areas of invasion have led to severe impacts on native flora and fauna. It is thought to have spread by means of released game fish bait and/or from aquarium release. Rusty crayfish are also raised for commercial and biological harvest.

Rusty crayfish reduce the amount and types of aquatic plants, invertebrate populations, and some fish populations--especially bluegill, smallmouth and largemouth bass, lake trout and walleye. They deprive native fish of their prey and cover and out-compete native crayfish. Rusty crayfish will also attack the feet of swimmers. On the positive side, rusty crayfish can be a food source for larger game fish and are commercially harvested for human consumption.

Rusty crayfish may be controlled by restoring predators like bass and sunfish populations. Preventing further introduction is important and may be accomplished by educating anglers, trappers, bait dealers and science teachers of their hazards. Use of chemical pesticides is an option, but does not target this species and will kill other aquatic organisms.

It is illegal to possess both live crayfish and angling equipment simultaneously on any inland Wisconsin water (except the Mississippi River). It is also illegal to release crayfish into a water of the state without a permit.




Figure 23 - Rusty Crayfish and identifying characteristics

ZEBRA MUSSELS

Zebra mussels (Figure 24) are an invasive species that have inhabited Wisconsin waters and are displacing native species, disrupting ecosystems, and affecting citizens' livelihoods and quality of life. They hamper

boating, swimming, fishing, hunting, hiking, and other recreation, and take an economic toll on commercial, agricultural, forestry, and aquacultural resources. The zebra mussel is a tiny (1/8-inch to 2-inch) bottom-dwelling clam native to Europe and Asia. Zebra mussels were introduced into the Great Lakes in 1985 or 1986, and have been spreading throughout them since that time. They were most likely brought to North America as larvae in ballast water of ships that traveled from fresh-water Eurasian ports to the Great Lakes. Zebra mussels look like small clams with a yellowish or brownish D-shaped shell, usually with alternating dark- and light-colored stripes. They can be up to two inches long, but most are under an inch. Zebra mussels usually grow in clusters containing numerous individuals.

Zebra mussels feed by drawing water into their bodies and filtering out most of the suspended microscopic plants, animals and debris for food. This process can lead to increased water clarity and a depleted food supply for other aquatic organisms, including fish. The higher light penetration fosters growth of rooted aquatic plants which, although creating more habitat for small fish, may inhibit the larger, predatory fish from finding their food. This thicker plant growth can also interfere with boaters, anglers and swimmers. Zebra mussel infestations may also promote the growth of blue-green algae, since they avoid consuming this type of algae but not others.

Zebra mussels attach to the shells of native mussels in great masses, effectively smothering them. A survey by the Army Corps of Engineers in the East Channel of the Mississippi River at Prairie du Chien revealed a substantial reduction in the diversity and density of native mussels due to Zebra Mussel infestations. The East Channel provides habitat for one of the best mussel beds in the Upper Mississippi River. Future efforts are being considered to relocate such native mussel beds to waters that are less likely to be impacted by zebra mussels.

Once zebra mussels are established in a water body, very little can be done to control them. It is therefore crucial to take all possible measures to prevent their introduction in the first place. Some of the preventative and physical control measures include physical removal, industrial vacuums, and back flushing.

Chemical applications include solutions of chlorine, bromine, potassium permanganate and even oxygen deprivation. An ozonation process is under investigation (patented by Bollyky Associates Inc.) which involves the pumping of high concentrations of dissolved ozone into the intake of raw water pipes. This method only works in controlling veligers, and supposedly has little negative impacts on the ecosystem. Further research on effective industrial control measures that minimize negative impacts on ecosystem health is needed.

In the fall of 2016, zebra mussels were found in a northwest Wisconsin lake for the first time. With this discovery, it increases the likelihood that zebra mussels will spread faster throughout northwest Wisconsin.

Zebra mussels have not been identified in Round Lake. Due to their incredibly invasive nature, a study was conducted several years ago to assess the suitability of lakes for zebra mussels. The result of that study was an on-line application called AIS Smart Prevention database which ranks lakes in WI as suitable, borderline suitable, or not suitable habitat for zebra mussel survival. This application uses statistical models of several variables to determine suitability rankings. In the fall of 2016, zebra mussels were first found in a lake in Northwestern Wisconsin that had been listed as suitable. Round Lake is listed as borderline suitable, so it is possible that a zebra mussel population could potentially become established if preventative measures are not taken.



Figure 24 - Zebra Mussels

AIS PREVENTION STRATEGY

Round Lake already has several established AIS. However there are many more that could be introduced to the lake. The RTLIA has and will continue to implement a watercraft inspection and AIS Signage program at the public access point on the lake. Information will be shared with lake residents and users in an effort to expand the watercraft inspection message. In addition to the watercraft inspection program, an in-lake and shoreland AIS monitoring program will be implemented. Both of these programs will follow UW-Extension Lakes and WDNR protocol through the Clean Boats, Clean Waters program and the Citizen Lake Monitoring Network AIS Monitoring program.

Additionally, having educated and informed lake residents is the best way to keep non-native AIS at bay in Round Lake. To foster this, the RTLIA will host and/or sponsor lake community events including AIS identification and management workshops; distribute education and information materials to lake property owners and lake users through the newsletter, webpage, and general mailings.

MANAGEMENT ALTERNATIVES

Nuisance aquatic plants can be managed a variety of ways in Wisconsin. The best management strategy will be different for each lake and depends on which nuisance species needs to be controlled, how widespread the problem is, and the other plants and wildlife in the lake. In many cases, an integrated pest management (IPM) approach to aquatic plant management that utilizes a number of control methods is necessary. The eradication of non-native aquatic invasive plant species such as EWM is generally not feasible, but preventing them from becoming a more significant problem is an attainable goal. It is important to remember however, that regardless of the plant species targeted for control, sometimes no manipulation of the aquatic plant community is the best management option. Plant management activities can be disruptive to a lake ecosystem and should not be done unless it can be shown they will be beneficial and occur with minimal negative ecological impacts.

Management alternatives for nuisance aquatic plants can be grouped into four broad categories: manual and mechanical removal, chemical application, biological control, and physical habitat alteration. Manual and mechanical removal methods include pulling, cutting, raking, harvesting, suction harvesting, and other means of removing the physical plant from the water and in most cases will require a WDNR permit. Chemical application is typified by the use of herbicides that kill or impede the growth of the aquatic plant and always requires a WDNR permit. Biological control methods include organisms that use the plant for a food source or parasitic organisms that use the plant as a host, killing or weakening it. Biological control may also include the use of species that compete successfully with the nuisance species for available resources. This activity may require a WDNR permit. Physical habitat alteration includes dredging, installing lake-bottom covers, manipulating light penetration, flooding, and drawdown. These activities may require permits under the WDNR waterways and wetlands program. It may also include making changes to or in the watershed of a body of water to reduce nutrients going in.

Each of the above control categories are regulated by the WDNR and most activities require a permit from the WDNR to implement. Mechanical harvesting of aquatic plants and under certain circumstances, physical removal of aquatic plants, is regulated under Wisconsin Administrative Rule NR 109 (Appendix A). The use of chemicals and biological controls are regulated under Administrative Rule NR 107 (Appendix B). Certain habitat altering techniques like the installation of bottom covers and dredging require a Chapter 30/31 waterway protection permit. In addition, anytime wild rice is involved one or more of these permits will be required.

Informed decision-making on aquatic plant management implementation requires an understanding of plant management alternatives and how appropriate and acceptable each alternative is for a given lake. The following sections list scientifically recognized and approved alternatives for controlling aquatic vegetation.

NO MANAGEMENT

When evaluating the various management techniques, the assumption is erroneously made that doing nothing is environmentally neutral. In dealing with nonnative species like CLP and EWM, the environmental consequences of doing nothing may be high, possibly even higher than any of the effects of management techniques. Unmanaged, these species can have severe negative effects on water quality, native plant distribution, abundance and diversity, and the abundance and diversity of aquatic insects and fish (Madsen, 1997). Nonindigenous aquatic plants are the problem, and the management techniques are the collective solution. Nonnative plants are a biological pollutant that increases geometrically, a pollutant with a very long residence time and the potential to "biomagnify" in lakes, rivers, and wetlands (Madsen, 2000).

Foregoing any management of AIS in Round Lake is not a recommended option. To keep EWM and CLP from causing greater harm, AIS management will continue to be implemented.

HAND-PULLING/MANUAL REMOVAL

Manual or physical removal of aquatic plants by means of a hand-held rake or cutting implement; or by pulling the plants from the lake bottom by hand is allowed by the WDNR without a permit per NR 109.06 Waivers under the following conditions:

- Removal of native plants is limited to a single area with a maximum width of no more than 30 feet measured along the shoreline provided that any piers, boatlifts, swim rafts and other recreational and water use devices are located within that 30-foot wide zone and may not be in a new area or additional to an area where plants are controlled by another method (Figure 25)
- Removal of nonnative or invasive aquatic plants as designated under s. NR 109.07 is performed in a manner that does not harm the native aquatic plant community
- Removal of dislodged aquatic plants that drift on-shore and accumulate along the waterfront is completed.
- The area of removal is not located in a sensitive area as defined by the department under s. NR 107.05 (3) (i) 1, or in an area known to contain threatened or endangered resources or floating bogs
- Removal does not interfere with the rights of other riparian owners
- If wild rice is involved, the procedures of s. NR 19.09 (1) are followed.




Figure 25 - Aquatic vegetation manual removal zone

Although up to 30 feet of aquatic vegetation can be removed, removal should only be done to the extent necessary. There is no limit as to how far out into the lake the 30-ft zone can extend, however clearing large swaths of aquatic plants not only disrupts lake habits, it also creates open areas for non-native species to establish. Physical removal of aquatic plants requires a permit if the removal area is located in a “sensitive” or critical habitat area previously designated by the WDNR. Manual or physical removal can be effective at controlling individual plants or small areas of plant growth. It limits disturbance to the lake bottom, is inexpensive, and can be practiced by many lake residents. In shallow, hard bottom areas of a lake, or where impacts to fish spawning habitat need to be minimized, this is the best form of control. If water clarity in a body of water is such that aquatic plants can be seen in deeper water, pulling AIS while snorkeling or scuba diving is also allowable without a permit according to the conditions in NR 106.06(2) and can be effective at slowing the spread of a new AIS infestation within a lake when done properly.

Larger scale hand or diver removal projects have had positive impacts in temporarily reducing or controlling AIS. Typically hand or diver removal is used when AIS has been newly identified and still exists as single plants or isolated small beds, but at least in one lake in New York State, it was used as a means to control a large-scale infestation of EWM. Kelting and Laxson (2010) reported that from 2004 to 2006 an “intensive

management effort” which involved “the selective removal of Eurasian water milfoil using diver hand harvesting of the entire littoral zone of the lake at least twice each summer for three years” followed by three years of maintenance management successfully reduced the overall distribution of EWM in the lake.

In Round Lake, there are some areas where this form of AIS control would be effective at helping control the spread of AIS. However, all of the AIS in Round Lake cannot be managed this way. Properly educating lake residents will allow them to remove AIS on and adjacent to their property.

DIVER ASSISTED SUCTION HARVESTING

Diver assisted suction harvesting or DASH, as it is often called, is a fairly recent aquatic plant removal technique. It is called "harvesting" rather than "dredging" because, although a specialized small-scale dredge is used, bottom sediment is not removed from the system. The operation involves hand-pulling of the target plants from the lake bed and inserting them into an underwater vacuum system that sucks up plants and their root systems taking them to the surface. It requires water pumps on the surface (generally on a pontoon system) to move a large volume of water to maintain adequate suction of materials that the divers are processing (Figure 26). Only clean water goes through the pump. The material placed by the divers into the suction hose along with the water is deposited into mesh bags on the surface with the water leaving through the holes in the bag. The bags have a large enough 'mesh' size so that silts, clay, leaves and other plant material being collected do not immediately clog them and block water movement. If a fish or other living marine life is sucked into the suction hose it comes out the discharge unharmed and is returned to the body of water. It can have some negative impacts to other nearby non-target plants if not done carefully, particularly those plants that are perennials and expand their populations by sub-sediment runners (Eichler, 1993).




Figure 26 - DASH - Diver Assisted Suction Harvest (Aquacleaner Environmental, <http://www.aquacleaner.com/index.html>); Many Waters, LLC)

In Wisconsin and Michigan, suction harvesting of invasive species is gaining popularity as a treatment method. There are several companies in the mid-west that are offering DASH services. Some of these companies are also building equipment that lake organizations and consultants can purchase to start up their own DASH program. Aquacleaner Environmental, out of Lancaster, NY sells a DASH system with a 5” suction hose for about \$30,000.00 plus extras. The same company offers DASH services at a rate of \$200.00/hour, with an acre of vegetation removal averaging \$15,000.00. Another company, Naturally DASH and Dredge, LLC (<http://www.naturallydash.com>), builds a system with a single pump and 3” hose for about \$6,000.00.

More locally, Many Waters, LLC out of Iron River, MI has been providing DASH services in northeastern WI. During the Northern Great Lakes Invasive Species Conference in Marquette, MI on November 4, 2014,

Many Waters, LLC presented DASH results from Lac Vieux Desert in Vilas County. During that presentation it was reported that 1,033.5 lbs. of EWM was removed from the lake with 17 hours of DASH. During the harvest, there was a 14.6% bi-catch of other plants sucked up at the same time. No report of costs was given. In this presentation, Many Waters, LLC reported that the efficiency of DASH was negatively impacted by obstacles/structures in the water, water clarity, sediment type, EWM density, native aquatic plant density, and time of year.

In a similar report filed for 2013 DASH services on Lake Elwood in Florence County, 2,322 lbs. of hybrid EWM was removed from the lake in 21 hours. In this lake, there was only a 1.85% bi-catch of native plants. According to documents on the Lake Ellwood Association webpage \$4,530.00 was spent on DASH services in 2013. Four areas in the lake totaling 0.7 acres were included in the DASH project. Based on these numbers, cost per lb. of EWM harvested was \$1.95; cost per hour for DASH services was \$215.71; and cost per acre was \$6,471.43. Lake Ellwood is a clear-water lake; however, the report mentions that DASH results were hampered by the presence of woody debris in the area of EWM harvest.

From a 2014 report for DASH services on Virgin Lake in Oneida County, 144 lbs. of EWM were removed in 2.5 hours with a bi-catch of 23%. On Virgin Lake, dense growth native vegetation and water clarity issues impacted the success of the DASH project. No report of cost was given.

More recently, June 2016, DASH was implemented on the St. Croix Flowage in Douglas County to remove approximately 2.0 acres of EWM, some dense, and some just scattered plants mixed in with many native plants. A new company TSB Lake Restoration was hired out of Chippewa Falls, WI for two days of DASH services. Approximately 16 hours of on the water time removed EWM from about 1.5 acres. The cost for the DASH services only was \$3,900.00 for the entire job. Broken down, the cost per acre was \$2,600.00; per hour was \$244.00; and per day was \$1,950.00. Consultant support costs added another \$1,800.00.

No formal documentation or measurement was made of lbs. of EWM removed or native species collateral damage, but observations by the Consultant estimate 500-800 lbs. of vegetation was removed and that up to about 30% of the plant material removed was non-target species, primarily common waterweed which dominated much of the bottom of the managed area. Collateral damage was the result of bringing the suction dredge too close to the bottom when feeding EWM into the tube.

MECHANICAL REMOVAL

Mechanical management involves the use of devices not solely powered by human means to aid removal. This includes gas and electric motors, ATV's, boats, tractors, etc. Using these instruments to pull, cut, grind, or rotovate aquatic plants is illegal in Wisconsin without a permit. DASH is also considered mechanical removal. To implement mechanical removal of aquatic plants a Mechanical/Manual Aquatic Plant Control Application is required annually. The application is reviewed by the WDNR and other entities and a permit awarded if required criteria are met. Using repeated mechanical disturbance such as bottom rollers or sweepers can be effective at control in small areas, but in Wisconsin these devices are illegal and generally not permitted.

LARGE-SCALE MECHANICAL HARVESTING

Large-scale mechanical harvesting is traditionally used for control of CLP, but can be an effective way to reduce EWM biomass in a water body. It is typically used to open up channels through existing beds of EWM to improve access for both human related activities like boating, and natural activities like fish distribution and mobility on lakes in maintenance mode where EWM is well-established and restoration efforts have been discontinued.

Aquatic plant harvesters are floating machines that cut and remove vegetation from the water. The size and harvesting capabilities of these machines vary greatly. As they move, harvesters cut a swath of aquatic plants

that is between 4 and 20 feet wide, and can be up to 10 feet deep. The on-board storage capacity of a harvester ranges from 100 to 1,000 cubic feet (by volume) or 1 to 8 tons (by weight). Most harvesters can cut between 2 and 8 acres of aquatic vegetation per day, and the average lifetime of a mechanical harvester is 10 years.

Mechanical harvesting of aquatic plants presents both positive and negative consequences to any lake. Its results - open water and accessible boat lanes - are immediate, and can be enjoyed without the restrictions on lake use which follow herbicide treatments. In addition to the human use benefits, the clearing of thick aquatic plant beds may also increase the growth and survival of some fish. By eliminating the upper canopy, harvesting reduces the shading caused by aquatic plants. The nutrients stored in the plants are also removed from the lake, and the sedimentation that would normally occur as a result of the decay of this plant matter is prevented. Additionally, repeated harvesting may result in thinner, more scattered growth.

Aside from the obvious effort and expense of harvesting aquatic plants, there are many environmentally-detrimental consequences to consider. The removal of aquatic species during harvesting is non-selective. Native and invasive species alike are removed from the target area. This loss of plants results in a subsequent loss of the functions they perform, including sediment stabilization and wave absorption. Shoreline erosion may therefore increase. Other organisms such as fish, reptiles, and insects are often displaced or removed from the lake in the harvesting process. This may have adverse effects on these organisms' populations as well as the lake ecosystem as a whole.

While the results of harvesting aquatic plants may be short term, the negative consequences are not so short lived. Much like mowing a lawn, harvesting must be conducted numerous times throughout the growing season. Although the harvester collects most of the plants that it cuts, some plant fragments inevitably persist in the water. This may allow the invasive plant species to propagate and colonize in new, previously unaffected areas of the lake. Harvesting may also result in re-suspension of contaminated sediments and the excess nutrients they contain.

Disposal sites are a key component when considering the mechanical harvesting of aquatic plants. The sites must be on shore and upland to make sure the plants and their reproductive structures don't make their way back into the lake or to other lakes. The number of available disposal sites and their distance from the targeted harvesting areas will determine the efficiency of the operation, in terms of time as well as cost.

Timing is also important. The ideal time to harvest, in order to maximize the efficiency of the harvester, is just before the aquatic plants break the surface of the lake. For CLP, it should also be before the plants form turions (reproductive structures) to avoid spreading the turions within the lake. If the harvesting work is contracted, the equipment should be inspected before and after it enters the lake. Since these machines travel from lake to lake, they may carry plant fragments with them, and facilitate the spread of AIS from one body of water to another. Harvesting contractors are not readily available in northern Wisconsin, so harvesting contracts are likely to be very expensive.

The extent of EWM and CLP in Round Lake does not warrant management of this scale which is why large-scale harvesting is not recommended.

SMALL-SCALE CUTTING WITH REMOVAL

There are a wide range of small-scale mechanical harvesting techniques, most of which involve the use of boat mounted rakes, scythes, and electric cutters. As with all mechanical harvesting, removing the cut plants is required. Commercial rakes and cutters range in prices from \$200 for rakes to around \$3000 for electric cutters with a wide range of sizes and capacities. Any of these cutters that are not entirely human powered require a permit for use.

Using a weed rake or cutter that is run by human power is allowed without a permit, but the use of any device that includes a motor, gas or electric, would require a permit. Dragging a bed spring or bar behind a boat, tractor or any other motorized vehicle to remove vegetation is also illegal without a permit. Although not truly considered mechanical management, incidental plant disruption by normal boat traffic is a legal method of management. Active use of an area is often one of the best ways for riparian owners to gain navigation relief near their docks. Most aquatic plants won't grow well in an area actively used for boating and swimming. It should be noted that purposefully navigating a boat to clear large areas is not only potentially illegal it can also re-suspend sediments, encourage AIS growth, and cause ecological disruptions.

Small-scale harvesting by human power that is completed in a way such that all of the EWM or CLP plant and root structure is removed is recommended for limited control of AIS in the lake. When removing aquatic plants manually, there is a restriction of no more than 30 feet wide for property owners to remove native vegetation, but there is no such limit on AIS. If done in a way that is entirely human powered, there is no limit on the amount of AIS that can be removed from the system. Any plants that are cut or pulled must still be removed from the lake. Through information and training, property owners will be instructed on proper physical removal methods.

BOTTOM BARRIERS AND SHADING

Physical barriers, fabric or other, placed on the bottom of the lake to reduce AIS growth would eliminate all plants, inhibit fish spawning, affect benthic invertebrates, and could cause anaerobic conditions which may release excess nutrients from the sediment. Gas build-up beneath these barriers can cause them to dislodge from the bottom and sediment can build up on them allowing AIS to re-establish. Bottom barriers are typically used for very small areas and provide only limited relief. Currently the WDNR does not permit this type of control.

Creating conditions in a lake that may serve to shade out EWM growth has also been tried with mixed success. The general intention is to reduce light penetration in the water which in turns limits the depth at which plants can grow. Typically dyes have been added to a small water body to darken the water. Bottom barriers and attempts to further reduce light penetration in Round Lake are not recommended.

DREDGING

Dredging is the removal of bottom sediment from a lake. Its success is based on altering the target plant's environment. It is not usually performed solely for aquatic plant management but rather to restore lakes that have been filled in with sediment, have excess nutrients, inadequate pelagic and hypolimnetic zones, need deepening, or require removal of toxic substances (Peterson, 1982). In shallow lakes with excess plant growth, dredging can make areas of the lake too deep for plant growth. It can also remove significant plant root structures, seeds/turions, rhizomes, tubers, etc. In Collins Lake, New York the biomass of curly-leaf pondweed remained significantly lower than pre-dredging levels 10-yrs after dredging (Tobiessen, Swart, & Benjamin, 1992). Dredging is very expensive, requires disposal of sediments, and has major environmental impacts. It is not a selective procedure so it can't be used to target any one particular species with great success except under extenuating circumstances. Dredging at any level must be permitted by the WDNR if done through mechanical means. Manual dredging of up to 100 cubic feet for may be done without a permit by property owners if all criteria found in the WDNR's exemption checklist (Appendix C) are met. It should not be performed for aquatic plant management alone. It is best used as a multipurpose lake remediation technique (Madsen, 2000).

Dredging is not a recommended management action for Round Lake.

DRAWDOWN

Drawdown, like dredging, alters the plant environment by removing all water in a water body to a certain depth, exposing bottom sediments to seasonal changes including temperature and precipitation. A winter

drawdown is a low cost and effective management tool for the long-term control of certain susceptible species of nuisance aquatic plants. Winter drawdown has been shown to be an effective control measure for EWM, but typically only provides 2-3 years of relief before EWM levels return to pre-drawdown levels. A winter drawdown controls susceptible aquatic plants by dewatering a portion of the lake bottom over the winter, and subsequently exposing vascular plants to the combined effect of freezing and desiccation (drying). The effectiveness of drawdown to control plants hinges on the combined effect of the freezing and drying. If freezing and dry conditions are not sustained for 4-6 weeks, the effectiveness of the drawdown may be diminished.

It is not possible to draw down Round Lake as there is no way to manipulate the water level at the existing outlet.

BIOLOGICAL CONTROL

Biological control involves using one plant, animal, or pathogen as a means to control a target species in the same environment. The goal of biological control is to weaken, reduce the spread, or eliminate the unwanted population so that native or more desirable populations can make a comeback. Care must be taken however, to insure that the control species does not become as big a problem as the one that is being controlled. A special permit is required in Wisconsin before any biological control measure can be introduced into a new area.

EWM WEEVILS

While many biological controls have been studied, only one has proven to be effective at controlling EWM under the right circumstances. *Euhrychiopsis lecontei* is an aquatic weevil native to Wisconsin that feeds on aquatic milfoils (Figure 28). Their host plant is typically northern watermilfoil; however they seem to prefer EWM when it is available. Milfoil weevils are typically present in low numbers wherever northern or Eurasian water milfoil is found. They often produce several generations in a given year and over winter in undisturbed shorelines around the lake. All aspects of the weevil's life cycle can affect the plant. Adults feed on the plant and lay their eggs. The eggs hatch and the larva feed on the plant. As the larva mature they eventually burrow into the stem of the plant. When they emerge as adults later, the hole left in the stem reduces buoyancy often causing the stem to collapse. The resulting interruption in the flow of carbohydrates to the root crowns reduces the plant's ability to store carbohydrates for over wintering reducing the health and vigor (Newman, Holmberg, Biesboer, & Penner, 1996).



Figure 27 - EWM Weevil (<https://klsa.wordpress.com/published-material/milfoil-weevil-guide/>)

The weevil is not a silver bullet. They do not work in all situations. The extent to which weevils exist naturally in a lake, adequate shore land over wintering habitat, the population of bluegills and sunfish in a system, and water quality characteristics are all factors that have been shown to affect the success rate of the weevil. The use of weevils is not recommended in this management plan, particularly since the process necessary to do so has changed significantly in the last few years. There is no longer a company that “raises” weevils for EWM

control. Weevils can still be raised by volunteers in cooperation with an overseeing entity, but requires that all EWM used in the rearing process be secured from the host lake, and only weevils reared on host lake EWM can be released into the host lake. Further monitoring and possible weevil rearing is not recommended for Round Lake in this management plan, but would not hurt if there were interested people to do so on the lake.

GALERUCELLA BEETLES

Two species of *Galerucella* beetles are currently approved for the control of purple loosestrife in Wisconsin (Figure 29). The entire lifecycle of *Galerucella* beetles is dependent on purple loosestrife. In the spring, adults emerge from the leaf litter below old loosestrife plants. The adults then begin to feed on the plant for several days until they begin to reproduce. Females lay their eggs on loosestrife leaves and stems. When the larvae emerge from these eggs they begin feeding on the leaves and developing shoots. When water levels are high these larvae will burrow into the loosestrife stems to pupate into adult beetles. These new adults emerge and begin feeding on the loosestrife again (Sebolt, 1998). *Galerucella* beetles do not forage on any other plants. Because of this the populations, once established, are self-regulating. When the purple loosestrife population drops off, the beetle population also declines. When the loosestrife returns, the beetle numbers will usually increase.



Figure 28 - Galerucella beetle

These beetles will not eradicate purple loosestrife entirely. This is true of almost all forms of biological control. *Galerucella* beetles will help regulate loosestrife which will allow native plants to also become established. This allows the wetlands surrounding Round Lake to be diverse plant communities instead of purple loosestrife monocultures.

Beetles can be obtained from the WDNR or at many of the public wetlands around Wisconsin. Because rearing these beetles requires the cultivation of a restricted species, a permit is necessary. Beetle rearing and release is not recommended for Round Lake in this management plan, but if lake residents wish to do so, it would not be detrimental.

OTHER BIOLOGICAL CONTROLS

There are other forms of biological control being used or researched. It was thought at one time that the introduction of plant eating carp could be successful. It has since been shown that these carp have a preference list for certain aquatic plants. EWM is very low on this preference list (Pine & Anderson, 1991). Use of “grass carp” as they are referred to in Wisconsin is illegal as there are many other environmental concerns including what happens once the target species is destroyed, removal of the carp from the system, impacts to other fish and aquatic plants, and preventing escapees into other lakes and rivers. Several pathogens or fungi are currently being researched that when introduced by themselves or in combination with

herbicide application can effectively control EWM and lower the concentration of chemical used or the time of exposure necessary to kill the plant (Sorsa, Nordheim, & Andrews, 1988). None of these have currently been approved for use in Wisconsin and are not recommended for use in Round Lake.

NATIVE PLANT RESTORATION

A healthy population of native plants might slow invasion or reinvasion of non-native aquatic plants. It should be the goal of every management plan to protect existing native plants and restore native plants after the invasive species has been controlled. In many cases, a propagule bank probably exists that will help restore native plant communities after the invasive species is controlled (Gettsinger, Turner, Madson, & Netherland, 1997). This is certainly the case in Round Lake where there are several native plants within the system, but the macrophyte population is comprised primarily of coontail and AIS. By doing native plant restorations, the plant diversity could be improved which could improve the system as a whole.

CHEMICAL CONTROL

Aquatic herbicides are granules or liquid chemicals specifically formulated for use in water to kill plants or cease plant growth. Herbicides approved for aquatic use by the U.S. Environmental Protection Agency (EPA) are considered compatible with the aquatic environment when used according to label directions. Some individual states, including Wisconsin, also impose additional constraints on herbicide use.

The Wisconsin Department of Natural Resources evaluates the benefits of using a particular chemical at a specific site vs. the risk to non-target organisms, including threatened or endangered species, and may stop or limit treatments to protect them. The Department frequently places conditions on a permit to require that a minimal amount of herbicide is needed and to reduce potential non-target effects, in accordance with best management practices for the species being controlled. For example, certain herbicide treatments are required by permit conditions to be in spring because they are more effective, require less herbicide and reduce harm to native plant species. Spring treatments also means that, in most cases, the herbicide will be degraded by the time peak recreation on the water starts.

The WDNR encourages minimal herbicide use by requiring a strategic Aquatic Plant Management (APM) Plan for management projects over 10 acres or 10% of the water body or any projects receiving state grants. WDNR also requires consideration of alternative management strategies and integrated management strategies on permit applications and in developing an APM plan, when funding invasive species prevention efforts, and by encouraging the use of best management practices when issuing a permit. The Department also supervises treatments, requires that adjacent landowners are notified of a treatment and are given an opportunity to request a public meeting if they want, requires that the water body is posted to notify the public of treatment and usage restrictions, and requires reporting after treatment occurs.

The advantages of using chemical herbicides for control of aquatic plant growth are the speed, ease and convenience of application, the relatively low cost, and the ability to somewhat selectively control particular plant types with certain herbicides. Disadvantages of using chemical herbicides include possible toxicity to aquatic animals or humans, oxygen depletion after plants die and decompose which can cause fishkills, a risk of increased algal blooms as nutrients are released into the water by the decaying plants, adverse effects on desirable aquatic plants, loss of fish habitat and food sources, water use restrictions, and a need to repeat treatments due to existing seed/turion banks and plant fragments. Chemical herbicide use can also create conditions favorable for non-native AIS to outcompete native plants (for example, areas of stressed native plants or devoid of plants).

When properly applied, the possible negative impacts of chemical herbicide use can be minimized. Early spring to early summer applications are preferred because exotic species are actively growing and many native plants are dormant, thus limiting the loss of desirable plant species; plant biomass is relatively low minimizing the impacts of de-oxygenation and contribution of organic matter to the sediments; fish spawning has ceased;

and recreational use is generally low limiting human contact. The concentration and amount of herbicides can be reduced because colder water temperatures enhance the herbicidal effects. Selectivity of herbicides can be increased with careful selection of application rates and seasonal timing. Lake hydro-dynamics must also be considered; steep drop-offs, inflowing waters, lake currents and wind can dilute chemical herbicides or increase herbicide drift and off-target injury. This is an especially important consideration when using herbicides near environmentally sensitive areas or where there may be conflicts with other water uses in the treatment vicinity.

Although done less frequently, herbicides can be applied in the late fall when most native plants have begun to die on their own, or have already gone dormant for the season. Typically invasive plant species like EWM will continue to grow well into the fall. Timing of a fall application of herbicides can be such that few native plants are expected to be killed. In some bodies of water, particularly those where wild rice is present, it may be possible to treat later in the fall, having no effect on wild rice that has already completed its life cycle. Wild rice in the seedling stage below the surface of the water is very susceptible to herbicides including 2, 4-D, endothall, and others. In most cases, herbicides are not used where wild rice is present. But in extreme cases, where the presence of EWM is actually causing great harm to the wild rice, fall treatments have been completed.

In some lakes, poor water clarity in the summer months may limit the growth of EWM, until the water clears in the fall and EWM all of a sudden gets more of the light needed to begin accelerated growth. The herbicide applied in the fall may be the same herbicide as applied in the spring and may be applied at the same concentration. One drawback is that the results of a fall treatment cannot be quantified until the next season.

HOW CHEMICAL CONTROL WORKS

Aquatic herbicides are sprayed directly onto floating or emergent aquatic plants or are applied to the water in either a liquid or granular form. Herbicides affect plants through either systemic or direct contact action. Systemic herbicides are capable of killing the entire plant. Contact herbicides cause the parts of the plant in contact with the herbicide to die back, leaving the roots alive and able to re-grow.

Herbicides can be classified as broad-spectrum (kill or injure a wide variety of plant species) or selective (effective on only certain species). Non-selective, broad spectrum herbicides will generally affect all plants that they come in contact with. Selective herbicides will affect only some plants. Often dicots, like Eurasian water milfoil, will be affected by selective herbicides whereas monocots, such as common waterweed will not be affected. The selectivity of a particular herbicide can be influenced by the method, timing, formulation, and concentration used.


Sonar® whose active ingredient is fluridone, is a broad spectrum herbicide that interferes with the necessary processes in a plant that create the chlorophyll needed to turn sunlight into plant food through a process called photo-synthesis. Rodeo® whose active ingredient is glyphosate is another broad spectrum herbicide that prevents an aquatic plant from making the protein it needs to grow. As a result the treated plant stops growing and eventually dies.

2, 4-D and triclopyr are active ingredients in several selective herbicides including Navigate®, DMA 4®, and Renovate®. These herbicides stimulate plant cell growth causing them to rupture, but primarily in dicots. These herbicides are considered selective as they have little to no effect on monocots in treated areas. Fluridone, glyphosate, 2, 4-D, and triclopyr are all considered systemic. When applied to the treatment area, plants in the treatment area draw the herbicide in through the leaves, stems, and roots killing all of the plant, not just the part that comes in contact with the herbicide.

Research done with triclopyr in 2014 (Vassios, et al., 2014) suggest that there is a difference between how the target plant is affected when using liquid or granular formulations of triclopyr. In short, liquid applications of

triclopyr tend to build up quicker in the meristem or growing tip of EWM, while granular applications tend to build up more in the root crown of EWM. The indication was that perhaps treating a body of water with both the granular and liquid formulation of the herbicide would affect a greater area of the plant providing better results than either formulation alone. This research was only completed using triclopyr, but it may have some application with 2,4-D as well, given that both herbicides affect the target plant in a similar way.

Aquathol whose active ingredient is endothall; Reward whose active ingredient is diquat; and Cutrine whose active ingredient is a form of copper are considered broad spectrum contact herbicides. They destroy the outer cell membrane of the material they come in contact with and therefore kill a plant very quickly. None of these three are considered selective and have the potential to kill all of the plant material that they come in contact with regardless of the species. As such, great care should be taken when using these products. Certain plant species like curly-leaf pondweed begin growing very early in the spring, even under the ice, and are often the only growing plant present at that time. This is a good time to use a contact herbicide like Aquathol, as few other plants would be impacted. Using these products later in the season, will kill all vegetation in contact with the herbicide and can provide substantial nuisance relief from a variety of aquatic plants.

It is possible to apply  more than one herbicide at a time when trying to establish control of unwanted aquatic vegetation. An example would be controlling EWM and CLP at the same time with an early season application, and controlling aquatic plants and algae at the same time during a mid-season nuisance relief application. Applying systemic and contact herbicides together has a synergistic effect leading to increased selectivity and control. Single applications of the two could result in reduced environmental loading of herbicides and monetary savings via a reduction in the overall amount of herbicide used and of the manpower and number of application periods required to complete the treatment.

EFFICACY OF AQUATIC HERBICIDES

The efficacy of aquatic herbicides is dependent on both application concentration and exposure time, and these factors are influenced by two separate but interconnected processes - dissipation and degradation. Dissipation is the physical movement of the active herbicide within the water column both vertically and horizontally. Dissipation rates are affected by wind, water flow, treatment area relative to untreated area, and water depths. Degradation is the physical breakdown of the herbicide into inert components. Depending on the herbicide utilized, degradation occurs over time either through microbial or photolytic processes.

MICRO AND SMALL-SCALE HERBICIDE APPLICATION

The determining factor in designating chemical treatments as micro or small-scale is the size of the area being treated. Small-scale herbicide application involves treating areas less than 10 acres in size. The dividing line between small-scale and micro treatments is not clearly defined, but is generally considered to be less than 3 acres. Small-scale chemical application is usually completed in the early season (April through May). Micro treatments are as well, but may be used as follow-up spot treatments after an early season application, or in instances where a new infestation has been identified in a lake with EWM already or in a completely new lake. Recent research related to micro and small-scale herbicide application generally shows that these types of treatment are less effective than larger scale treatments due to rapid dilution and dispersion of the herbicide applied.

Some suggested ways to increase the effectiveness of this management strategy are to increase the concentration of herbicide used, use a contact herbicide like diquat that does not require as long a contact time to be effective, or in some manner contain the herbicide in the treated area by artificial means.

Pre- and post-treatment aquatic plant surveys and testing for herbicide residuals are not required by the WDNR for small-scale treatments. Nor is an approved APMP if the organization sponsoring the application is not using grant funding to help defer the costs. Even though not required by the WDNR, participating in

these activities is recommended as it helps to gain a better understanding of the impact and fate of the chemical used.

LARGE-SCALE HERBICIDE APPLICATION

Large-scale herbicide application involves treating areas more than 10 acres in size. Like small-scale applications, this is usually completed in the early-season (April through May) for control of non-native invasive species like EWM while minimizing impacts on native species. It is generally accepted that lower concentration of herbicide can be used in large-scale applications as the likelihood of the herbicide staying in contact with the target plant for a longer time is greater. If the volume of water treated is more than 10% of the volume of the lake, or the treatment area is ≥ 160 acres, or 50% of the lakes littoral zone, effects can be expected at a whole-lake scale. Large-scale herbicide application can be extended in some lakes to include whole bay or even whole lake treatments. The bigger the treatment area, the more contained the treatment area, and the depth of the water in the treatment area, are factors that impact how whole bay or whole lake treatments are implemented.

Pre- and post-treatment aquatic plant surveying and having an approved APMP are required by the WDNR when completing large-scale chemical treatments. Residual testing is not required by the WDNR, but highly recommended to gain a better understanding of the impact and fate of the chemical used.


WHOLE-LAKE, AND/OR EPILIMNION APPLICATION

Whole-lake or whole-basin treatments are those where the herbicide may be applied to specific sites, but the goal of the strategy is for the herbicide to reach a target concentration when it equally distributes throughout the entire volume of the lake (or lake basin, or within the epilimnion of the lake or lake basin). The application rate of whole-lake treatments is dictated by the volume of water in with which the herbicide will reach equilibrium. Because exposure time is expected to be so much longer, effective herbicide concentrations for whole-lake treatments are significantly less than required for spot treatments. Whole-lake treatments are typically conducted when the target plant is spread throughout the majority of the lake or basin.

If the herbicide exposure time of the target aquatic plant can be extended, the concentration of the herbicide applied can be lowered. If the contact time between the applied herbicide and the target plant in a whole body of water or protected bay can be increased to, or is already expected to be several days to a week or more, the concentration of herbicide can be in the range of 0.25-0.5 ppm instead of the 2-4 or more ppm that is typically used in small-scale, spot, or micro treatments.

Planning to treat the whole lake can be further designed to minimize the herbicide needed to affect the desired outcome. The method used to implement whole-lake treatments changes with the type of lake. Herbicide applied to a shallow, mixed lake is expected to mix throughout the entire volume of the lake. In deep water lakes that stratify, herbicide can be applied at such a time when it is expected that it will only mix with the surface water above the thermocline in an area known as the epilimnion (Figure 30). For this to be a viable management alternative, a lake has to stratify early enough in the open water season to coincide with the optimal time for early season chemical application.

Lakewide Vertical Dissipation



Slide Credit: Nault, Michelle and John Skogerboe. 2014. *Scientific Evaluation of Efficacy and Selectivity of Herbicide Treatments in Wisconsin Lakes*. Presentation at UMISC – October 22, 2014 in Duluth, Minnesota.

Figure 29 - Lake-wide (whole-lake) dissipation of aquatic herbicides in Mixed and Stratified Lakes (Carlson, 2015). Inset: Summer thermal stratification.

EFFECTS OF WHOLE-LAKE TREATMENTS ON NATIVE AQUATIC PLANT SPECIES

Treating an entire lake with a chemical herbicide does have some concerns. One is particular is the effect on native aquatic plant vegetation in the treated body of water. Based on study results published by the WDNR in 2012 (Nault, et al., 2012) looking at nine different lakes that had whole-lake treatments completed, “year of treatment” effects on native plants were mostly negative, and on aggregate, 34 of the total 38 significant differences between species frequency of occurrence pre- and post-treatment were reductions, affecting 38 percent to 78 percent of the number of native species within a lake. Short-term reductions in native littoral frequency of occurrence occurred even at low concentrations of 2, 4-D if exposure times were long. Native dicots such as the watermilfoils (esp. northern watermilfoil), water marigold, and bladderworts are known to be susceptible to 2, 4-D, and displayed statistically-significant decreases in some of the case studies. At long-term exposures (across a range of concentrations) adverse impacts to relatively tolerant monocots such as naiads, several narrow leaf pondweeds, wild celery, and common waterweed were also observed. Water quality may also be affected by large-scale treatments. For example, in two lakes for which Secchi data was collected pre- and post-treatment (Sandbar and Tomahawk), a 40-percent reduction in water clarity was observed when comparing pre-treatment averages to year-of treatment averages. In another Wisconsin lake not part of this study (Bridge Lake), dissolved oxygen levels declined following a large-scale treatment that occurred relatively late in the season when water temperatures were higher.

PRE AND POST TREATMENT AQUATIC PLANT SURVEYING

When introducing new chemical treatments to lakes where the treatment size is greater than ten acres or greater than 10% of the lake littoral area and more than 150-ft from shore, the WDNR requires pre and post chemical application aquatic plant surveying. The protocol for pre and post treatment survey is applicable for chemical treatment of CLP and EWM.

The WDNR protocol assumes that an APMP has identified specific goals for non-native invasive species and native plants species. Such goals could include reducing coverage by a certain percent, reducing treatments to

below large-scale application designations, and/or reducing density from one level to a lower level. A native plant goal might be to see no significant negative change in native plant diversity, distribution, or density. Results from pre and post treatment surveying are used to improve consistency in analysis and reporting, and in making the next season's management recommendations.

The number of pre and post treatment sampling points required is based on the size of the treatment area. Ten to twenty acres generally requires at least 100 sample points. Thirty to forty acres requires at least 120 to 160 sampling points. Areas larger than 40 acres may require as many as 200 to 400 sampling points. Regardless of the number of points, each designated point is sampled by rake, recording depth, substrate type, and the identity and density of each plant pulled out, native or invasive.

In the year prior to an actual treatment, the area to be treated must have a mid-season/summer/warm water point intercept survey completed that identifies the target plant and other plant species that are present. A pre-treatment aquatic plant survey is done in the year the herbicide is to be applied, prior to application to confirm the presence and level of growth of the target species. A post-treatment survey should be scheduled when native plants are well established, generally mid-July through mid-August. For the post-treatment survey, repeat the PI for all species in the treatment polygons, as was done the previous summer. For whole-lake scale treatments, a full lake-wide PI survey should be conducted.

CHEMICAL CONCENTRATION TESTING

Chemical concentration testing is often done in conjunction with treatment to track the fate of the chemical herbicide used. Testing is completed to determine if target concentrations are met, to see if the chemical moved outside its expected zone, and to determine if the chemical breaks down in the system as expected. Monitoring sites are located both within and outside of the treatment area, particularly in areas that may be sensitive to the herbicide used, where chemical drift may have adverse impacts, where movement of water or some other characteristic may impact the effect of the chemical, and where there may be impacts to drinking and irrigation water. Water samples are collected prior to treatment and for a period of hours and/or days following chemical application.

In some lakes, rhodamine dye is added to the herbicide at the time of application in amounts equal to the expected concentration of the herbicide and a fluorimeter is used to sample the dye as it moves around the system. Both systems for tracking the movement of the herbicide, concentration attained, and contact time maintained can be used effectively to help better current and future planning.

HERBICIDE USE IN ROUND LAKE

Round Lake has a fairly limited native plant population. The level of EWM in the system creates an additional challenge for native plants within Round Lake. To minimize EWM's impact on the lake's native plants, every effort should be made to maintain it at or further reduce it from its current low levels. In the previous APM Plan, curly-leaf pondweed was not thought to be negatively impacting the lake.

Past chemical management has been completed using either a granular or liquid version of a commercial herbicide with 2, 4-D as the active ingredient. Navigate, a granular formulation with 2, 4-D is a common aquatic herbicide used for control of submerged aquatic plants. DMA 4, a liquid formulation with 2, 4-D is less expensive than Navigate but essentially is applied at the same concentrations that are used with a granular herbicide. Both of these commercial brands are considered systemic herbicides, expected to kill all parts of the plant, not just what it comes in contact with.

Triclopyr (Trade name Renovate) is another systemic herbicide approved for use in WI to control submersed aquatic vegetation like EWM. It too comes in granular or liquid formulations, and could be used instead of 2, 4-D based herbicides at comparable concentrations. Presently triclopyr based herbicides are more expensive than 2, 4-D based herbicides.

Research done with triclopyr in 2014 (Vassios, Nissen, Koschnick, & Heilman, 2014) suggest that there is a difference between how the target plant is affected when using liquid or granular formulations of triclopyr. In short, liquid applications of triclopyr tend to build up quicker in the meristem or growing tip of EWM, while granular applications tend to build up more in the root crown of EWM. The indication was that perhaps treating a body of water with both the granular and liquid formulation of the herbicide would affect a greater area of the plant providing better results than either formulation alone. This research was only completed using triclopyr, but it may have some application with 2,4-D as well, and it would be interesting to complete a test treatment using this method.

Diquat (Trade name Reward) was used in 2016, but the application of this herbicide had little impact on any of the EWM it was applied to, and the treatment was considered a failure. The use of 2,4-D based herbicides (granular or liquid) had mixed success primarily due to the amount of water that flows through Round Lake via the Trade River. Low or normal water levels are a must to consider chemical treatment of EWM or CLP in Round Lake in any given year. In addition, consideration of using 2,4-D and endothall in combination to treat CLP and EWM is recommended; or as an alternative, to apply a mix of both granular and liquid 2,4-D to potentially increase success rates. This latter idea, could be supported with a study where comparisons are made between areas treated with one or the other form of herbicide and areas treated with both.

Procellacor® is a new herbicide that acts similar to both contact and systemic herbicides. This allows for it to be effective with a shorter amount of contact time. The active ingredient in Procellacor® is an organic compound which mimics the plant hormone auxin. The auxins that are produced naturally within plants stimulate stem elongation and while suppressing bud growth. However when auxin concentrations within plant tissues reach a certain threshold, the growth response is completely reversed. The plant begins to, essentially, prepare for a dormant period by stopping growth altogether and abscising leaves. At this point, additional auxins (or their mimics) will become toxic to the plant and result in cell death. This herbicide is not currently approved for use within Wisconsin, but it is currently under review by through the Wisconsin Environmental Policy Act (WEPA). There is little information on the recommended time-frame or concentrations of treatments with Procellacor® due to the lack of State approval; however if approved for use within Wisconsin, this herbicide should be reviewed and considered for use in the Trade River system.

In order to effectively manage both EWM and CLP, both systemic and contact herbicides should be applied early in the season. This will allow EWM and CLP to be heavily impacted while native plants, which have not yet begun to grow, will be minimally affected.

MANAGEMENT DISCUSSION

EWM

The littoral (plant growing) zone of Round Lake in 2016 was approximately 40 acres, approximately 20% of the total surface area (208 acres). Since 2011, the amount of EWM in Round Lake as identified by fall bed-mapping surveys has fluctuated between 1.41 and 17.01 acres in beds (>50% EWM with a defined edge) and/or high density areas (>25% EWM with a defined edge) (Table 7). These values represent 3.3 to 42.5% of the littoral zone. The average amount of EWM during the fall surveys in years when herbicide treatments were considered more effective (2013-2015) was 2.75 acres. Based on these numbers it is a reasonable goal to keep the level of EWM in Round Lake as identified in a fall bed mapping survey, below 2.75 acres of the littoral zone (6.9%) in any given year.

Table 7 - EWM Distribution based on Fall Bed-mapping Surveys

Fall EWM Bed Mapping Results - Round Lake		
Year	Total Acres	Notes
2011	17.01	No treatment
2012	8.84	Treated with granular 2,4-D
2013	1.41	Treated with liquid and granular 2,4-
2014	3.2	Treated with liquid and granular 2,4-
2015	3.65	Treated with liquid and granular 2,4-
2016	7.54	Treated with diquat
2017	2.76	Treated with liquid 2,4-D

CLP

CLP is not as well established in Round Lake as it is in the other lakes, although its distribution and density seems to be increasing. Over the next five years, management on Round Lake will strive to control most of the EWM, and at the same time target CLP when present in the same places EWM is being targeted. Reducing the amount of moderate to dense growth CLP by 50% in five years would lessen the undesirable impacts of the plant in the lake which include interfering with native aquatic plant growth in the spring, causing navigation and nuisance conditions in parts of the lake in the late spring and early summer, and contributing to nutrient loading and organic material build up in the sediment mid-summer..

APPLICATION OF AQUATIC HERBICIDES

If the total amount of EWM identified during a fall bed-mapping survey of the littoral zone reaches or exceeds 2.75 acres then the application of herbicide will be considered. Proposed treatment areas will not be less than 1.0 acres in size. Any bed or high density area of EWM that exceeds 0.02 acres (approximately 900-ft²) will be included in a preliminary early season treatment proposal for the given year. A buffer of 25-50 feet will be established around identified beds. Gaps between beds may also be included in proposed treatment areas if the two beds in question are close to one another, or if the area between the two beds has been known to support EWM growth based on past mapping actions. If herbicides are incorporated in a treatment plan for a given year, both CLP and EWM will be targeted at the same time using endothall at 1.0-3.0 ppm and 2,4-D at 2.0-4.0 ppm. Assuming treatment areas are at least one acre in size, liquid formulations of these herbicides will be used.

CLP reductions will be accomplished by treating it along with any EWM treatment that is implemented. Endothall and 2,4-D will be applied back to back. Previous results in Little Trade Lake (2013 & 2016) and Long Trade Lake (2013 & 2016) have shown this approach to be highly successful at controlling both species, for multiple years. Should all combined treatment areas exceed 10% (4 acres) of the littoral zone official pre and post treatment, point-intercept, aquatic plant surveys will be completed. Once CLP has been treated in an area for the first time, it will be continued for a minimum of two years more.

The success of these treatments will be measured by annual spring CLP bed-mapping, and by comparing turion density counts in the first year of treatment in this new plan (2018) with counts taken after a minimum of three years' worth of active management.

AQUATIC PLANT SURVEYING

Beginning in 2018, following year EWM treatment proposals will be based on late summer or early fall point-intercept survey work in the entire littoral zone of the lake. All aquatic plants in the littoral zone will be identified. If areas of EWM identified in beds and/or high density areas exceed 3.0 acres of the littoral zone or are causing a navigational impairment, a chemical treatment proposal will be made. If the size of the proposed treatment area reaches or exceeds 10% (4 acres) of the littoral zone, pre and post treatment point-intercept aquatic plant surveys will be completed within the treated areas. If the proposed treatment area is less than 10% of the littoral zone, pre and post treatment point-intercept aquatic plant surveys are not required but will be considered if resources are available to support it. If pre and post treatment survey work is not done, a EWM Readiness survey will be completed in the proposed treatment areas prior to actual treatment to determine if an appropriate amount and level of CLP and EWM growth has been attained to implement the treatment. After any survey work, modifications to the initial treatment proposal will be made if necessary. The late summer/early fall point-intercept survey of the entire littoral zone will be used to make annual comparisons of treatment impacts on target and non-target aquatic plant species from year to year.

OTHER AIS MONITORING AND MANAGEMENT

RTLIA volunteers will continue to monitor the shoreline for purple loosestrife, removing what is found if possible. The RTLIA will be involved in rearing beetles for biological control of purple loosestrife however where those beetles are released each year will be determined by the location and most dense areas of purple loosestrife. For at least the last 5 years biological control beetles have been released on Round Lake, as the other lakes to date do not have large areas of purple loosestrife.

No formally recognized management of reed canary grass or Chinese mystery snails is expected, although shoreland improvement projects completed during the time span of this plan might impact the level of reed canary grass along the shore.

RTLIA volunteers will participate in the CLMN AIS Monitoring Program annually looking for zebra mussels, rusty crayfish, hydrilla, and other AIS not already in the lake.

COARSE WOODY HABITAT

Coarse woody habitat has never been quantified in Round Lake. At some point during the implementation of this 5-year plan, the amount of CWH will be quantified and willing property owners sought for the installation of one or more CWH projects. Increasing the level of CWH in the lake would likely improve the overall fishery in the lake.

SHORELAND IMPROVEMENT

As increasing nutrients and sediment to the lake is a concern and has led to Round Lake being placed on the EPA/State of Wisconsin Impaired Waters list. Making improvements to the nearshore area around the lake and upstream of the lake in the Trade River could benefit the lake.

To that end, the WDNR has begun implementing a new Lake Shoreland and Shallows Habitat Monitoring Field Protocol (Appendix D) that involves assessing a 35-ft buffer area around the entire lake, documents shoreland condition through digital photography, and documents coarse woody debris in a lake. Information about the condition of the shoreline of Round Lake would benefit future shoreland improvement planning and implementation through the WDNR Healthy Lakes grant program and other programs sponsored by the RTLIA and Burnett County. Polk County completed a Shoreland Habitat Assessment project on Long Trade Lake in 2017. A similar approach could be done by Burnett County on Round Lake, although it is more likely that such an assessment would be completed by a consultant retained by the RTLIA with the cost being covered in a future lake management planning grant funded project aimed at developing a water quality based management plan for the lake. It is recommended that a shoreland habitat assessment survey be completed following the new WDNR protocol during the time frame covered by this APM Plan.

WATER QUALITY

It is recommended that the RTLIA continue to seek and support volunteers on Round Lake who collect basic water quality data through the Citizen Lake Monitoring Network (CLMN) water quality monitoring program. This program begins with the collection of water clarity data, but can be expanded to include temperature and oxygen profiling, and collection of water samples to be analyzed for total phosphorus and chlorophyll a. It would also be beneficial to determine water and nutrient budgets for the lake.

A water budget is an accounting of all the water that flows into and out of a project area. This area can be a wetland, a lake, or any other point of interest. Development can alter the natural supply of water and severely impact an area, especially if there are nearby ponds or wetlands. A water budget is needed to determine the magnitude of these impacts and to evaluate possible mitigation actions. Components of a water budget include: precipitation, evaporation, evapotranspiration, surface runoff, and groundwater flow. The first three terms of the water budget equation, precipitation, evaporation, and evapotranspiration, are natural processes that are largely unaffected by development. However, changes in land use can significantly affect surface runoff and groundwater flow. For example, commercial development may intercept surface runoff that ran into a wetland and redirect it to a stormwater control basin. This stormwater basin may hold the water until it evaporates or release it to an outlet stream. In either case, the wetland is deprived of the surface runoff that was available before the development. Similarly, water supply wells can permanently lower groundwater levels and change flow directions.

The calculation of a nutrient budget is an essential step in the evaluation of a lake's trophic status. A nutrient budget provides a means to evaluate and rank nutrient sources that may contribute to algal problems. It is most important to determine the quantity of nutrients (especially phosphorus) entering the lake, as well as the ultimate fate of those nutrients. Components of a nutrient budget include: tributary loading and discharge; atmospheric loading; direct surface runoff; septic leachate and groundwater loading; groundwater phosphorus, and internal release of phosphorus from the sediment of the lake.

2018-2022 AQUATIC PLANT MANAGEMENT GOALS, OBJECTIVES, AND ACTIONS

Based on the information shared in this document the following aquatic plant management goals, objectives, and actions are recommended. The Goals, Objectives, and Actions are also available in Appendix E.

Goal 1 – Promote and support aquatic plant management strategies that will control the spread of aquatic invasive species without negatively impacting native vegetation in Round Lake.

- 1) Objective 1 – Keep level of EWM to below 2.75 acres as indicated by annual summer littoral point-intercept surveys of aquatic vegetation
 - a) Action – Early season small-scale herbicide application
 - i) Treated areas must be ≥ 1 acre
 - ii) Combine EWM and CLP treatments if in the same proposed area
 - iii) Herbicide concentration/dispersion monitoring within the treated area and at the outlet of the lake
 - b) Action – Physical Removal of EWM
 - i) Summer littoral visual surveys with removal
 - ii) Property owner removal near docks
 - c) Action – EWM surveys
 - i) Mid to late summer PI aquatic plant surveys within the littoral zone
 - ii) Treatment readiness survey – mid-April to mid-May
 - iii) Pre- and post-treatment aquatic plant survey work
 - (1) Only implemented if proposed treatment areas exceed 10 acres
- 2) Objective 2 – Minimize negative impacts caused by dense growth CLP
 - a) Action – Early season small-scale herbicide application (w/EWM)
 - i) Treated areas must be ≥ 1 acres
 - ii) Combine EWM and CLP treatments if in the same proposed area
 - iii) Herbicide concentration/dispersion monitoring within the treated area and at the outlet of the lake
 - iv) Once implemented, must be completed for a minimum of three years
 - b) Action – CLP surveys
 - i) CLP bed-mapping – June
 - ii) Treatment readiness survey – mid-April to mid-May
 - iii) Pre- and post-treatment aquatic plant survey work
 - (1) Only implemented if proposed treatment areas exceed 10 acres
 - iv) Fall turion density sampling – 2018, and again after three years of active management
- 3) Objective 3 – Annual Summer Aquatic Plant Surveying
 - a) Action – Summer Littoral Point-Intercept Survey
 - i) All plants, entire littoral zone, point-intercept survey
 - ii) Mid-July to mid-August
 - iii) Year to year comparisons

Goal 2 – Reduce the threats that existing AIS will leave the lake; that new aquatic invasive species will be introduced into the lake; and that new AIS introduced to the lake will go undetected in the lake.

- 1) Objective 1 – Clean Boats, Clean Waters

- a) Action - 200 hours annually with grant funding
- b) Action - Volunteer hours only without grant funding
- 2) Objective 2 – AIS Monitoring
 - a) Action – Participate in CLMN AIS monitoring
 - b) Action – Implement fall dock and boatlift zebra mussel survey
 - c) Action – Maintain and/or improve AIS signage at landing
- 3) Objective 3 – AIS Education
 - a) Action – Distribute AIS education and identification materials
 - b) Action – Plan and implement AIS identification and physical removal workshops
- 4) Objective 4 – AIS Control
 - a) Action – Implement physical removal or other approved management techniques when necessary

Goal 3 – Promote and support nearshore and riparian best management practices that will improve fish and wildlife habitat, reduce runoff, and minimize nutrient loading into Round Lake.

- 1) Objective 1 – Implement State of Wisconsin Healthy Lakes Initiative
 - a) Action – Complete a Shoreland Habitat Assessment on Round Lake
 - b) Action - Promote Healthy Lakes projects based on the results of the Shoreland Habitat Assessment
 - c) Action – Apply for Healthy Lake grant funding to support projects that improve shoreland habitat and reduce runoff into the lake
- 2) Objective 2 – Direct attention to water quality issues and how they could be addressed
 - a) Action - Research existing water quality, watershed, and nearshore condition of the lake
 - b) Action – Determine what information is needed to move water quality planning forward
 - c) Action – Consider application for lake management planning grant funding to support initial water quality management data collection and planning.
 - i) Include a feasibility study to evaluate hydraulic and nutrient loading to the lake
 - ii) Complete in-lake monitoring to determine recycling and profile characteristics

Goal 4 – Complete appropriate and on-going tracking, monitoring, and management strategy modification to allow for thorough evaluation of management actions, and determinations that those management actions are on target, on track, on schedule, on budget, and within expected parameters.

- 1) Objective 1 – Continue water quality testing for Secchi, temperature, dissolved oxygen, total phosphorus, and chlorophyll a at the Deep Hole in Round Lake
 - a) Action – Continue involvement in the Citizen Lake Monitoring Network (CLMN)
 - b) Action – Purchase a Temp/DO meter to support water quality testing on Round Lake
- 2) Objective 2 – Complete Annual Project Activity and Assessment Reports
 - a) Action – The RTLIA and their Consultant will prepare end-of-year reports summarizes the management actions taken and how they impacted the lake.
 - b) Action – Review end of year summary reports with the RTLIA and WDNR to determine following year management actions.

Goal 5 – Encourage and engage lake residents and visitors to be active lake stewards.

- 1) Objective 1 – Promulgate behavior change in residents in the following areas: AIS, shoreland development, aquatic vegetation, and responsibility for the lake.
 - a) Action – Encourage lake residents to understand AIS concerns, learn to identify AIS, watch for and identify AIS in the lake, and report what they find and/or remove it
 - b) Action – Encourage boaters to implement appropriate AIS prevention strategies on their watercraft; observe no-wake rules for boats and PWC close to shore and to each other; and be considerate of others on the lake
 - c) Action – Encourage lake residents to let vegetation in the water grow and to plant native plants along their shore
 - d) Action – Encourage lake residents to care for their lake, not just their lawn
 - i) Provide education materials, welcome packets, newsletters, information/education displays, Facebook, webpage, meetings and other resources to increase the level of public awareness on the lake
 - ii) Establish and develop volunteer lake leadership
 - (1) Attend conferences
 - (2) Recruit new representatives from Round Lake to be active board/committee members
 - (3) Encourage more membership by Round Lake residents in the RTLIA
 - (4) Encourage lake volunteer involvement in “lake leaders” training
 - iii) Strive to engage the youth in preserving the future health of the lake
 - iv) Highlight examples of good shoreland practices on the lake
 - v) Recognize good lake stewards for the efforts they extend

Goal 6 – Implement the Round Lake Management Plan effectively and efficiently with a focus on community and constituent education, information, and involvement.

- 1) Objective 1 - Build and support partnerships.
 - a) Action – Work with WDNR, Burnett County, Town of Trade Lake, local businesses, contractors, and other resources to support management actions
- 2) Objective 2 – Keep lake residents are informed about plan activities
 - a) Action – Continue supporting Round Lake involvement in the RTLIA
 - b) Action – Continue reaching out to the lake constituency to inform and seek input for management actions
- 3) Objective 3 - Select cost effective implementation actions
 - a) Action – Work within the budget constraints to establish the best management actions to implement annually
 - b) Action – Apply for State of Wisconsin grant funding to support education, planning, and management implementation

IMPLEMENTATION AND EVALUATION

This plan is intended to be a tool for use by the RTLIA to move forward with aquatic plant management actions that will maintain the health and diversity of Round Lake and its aquatic plant community. This plan is not intended to be a static document, but rather a living document that will be evaluated on an annual basis and updated as necessary to ensure goals and community expectations are being met. This plan is also not intended to be put up on a shelf and ignored. Implementation of the actions in this plan through funding obtained from the WDNR and/or RTLIA funds is highly recommended. An Implementation and Funding Matrix is provided in Appendix F.

Since many actions occur annually, a calendar of actions to be implemented was created in Appendix G.

WISCONSIN DEPARTMENT OF NATURAL RESOURCES GRANT PROGRAMS

There are several WDNR grant programs that may be able to assist the SLMD in implementing its new APMP. AIS grants are specific to actions that involve education, prevention, planning, and in some cases implementation of AIS management actions. Lake Management Planning grants can be used to support a broad range of management planning and education actions. Lake Protection grants can be used to help implement approved management actions that would help to improve water quality. WDNR Healthy Lakes grants are part of the Lake Protection program.

AIS PREVENTION AND CONTROL GRANTS

The AIS (AIS) Prevention and Control grants are a cost-share effort by the WDNR to provide information and education on types of existing and potential AIS in Wisconsin, the threats that invasive species pose to the state's aquatic resources, and available techniques for invasive species control. These grants also assist in the planning and implementation of projects that will prevent the introduction of invasive species into waters where they currently are not present, controlling and reducing the spread of invasive species from waters where they are present, and restoring native aquatic communities.

There are five AIS Prevention and Control grants subprograms:

- Education, Prevention and Planning Projects (including Clean Boats Clean Waters)
- Early Detection and Response Projects
- Established Population Control Projects
- Maintenance and Containment Projects
- Research and Demonstration Projects

Education, Prevention, and Planning; Clean Boats, Clean Waters, and Maintenance and Containment grants are applicable to Round Lake and the RTLIA.

EDUCATION, PREVENTION AND PLANNING PROJECTS

Education projects are intended to broaden the public's awareness and understanding of, and ability to identify, AIS; the threats that AIS pose to the health of aquatic ecosystems; the measures to prevent the spread of AIS; and the management practices used for control of AIS. Prevention projects are intended to prevent the introduction of new AIS into a waterbody/wetland, or prevent the spread of an AIS population from one waterbody to another unpopulated waterbody/wetland. Planning projects are intended to assist in the development of plans for the prevention and control of AIS. Eligible projects include:

- Educational programs including workshops, training sessions, or coordinated volunteer monitors. Projects will be reviewed for consistency with the DNR's statewide education strategy for controlling AIS including the use of existing publications and outreach materials.
- Development of AIS prevention and control plans
- Monitoring, mapping, and assessing waterbodies for the presence of AIS or other studies that will aid in the AIS prevention and control.
- Watercraft inspection and education projects following the guidelines of the DNR's Clean Boats, Clean Waters program.

This subprogram is not intended to provide support for any management action that may be taken.

Funding Possibilities

Maximum amount of grant funding is 75% of the total project costs, not to exceed \$150,000. Applications will be separated into two classes: less than \$50,000 in state funding and between \$50,001 and \$150,000 in state funding. Clean Boats Clean Waters projects are limited to \$4,000 per public boat launch facility but may be a component of a larger project.

ESTABLISHED POPULATION CONTROL PROJECTS

Established population control grants are intended to assist applicants in eradicating or substantially reducing established populations of AIS to protect and restore native species communities. Established populations are defined as substantial reproducing populations of AIS that are not pioneer populations. Eligible projects include activities recommended in a DNR-approved control plan including monitoring, education, and prevention activities. Ineligible projects include the following:

- Dredging
- Chemical treatments or mechanical harvesting of aquatic plants to provide single season nuisance or navigational relief.
- Maintenance and operation of aeration systems and mechanical structures used to suppress aquatic plant growth.
- Structural facilities for providing boat washing stations. Equipment associated with boat washing facilities is eligible if included in a management plan.

Funding Possibilities

Maximum amount of the grant funding is 75% of the total project costs, not to exceed \$200,000.

MAINTENANCE AND CONTAINMENT PROJECTS

Maintenance and containment grants are intended to provide sponsors limited financial assistance for the ongoing control of established AIS population without the assistance of an Established Population Control grant. These projects are intended for waters where management activity has achieved the target level of control identified in an approved plan that meets the criteria of s. NR 198.43, Wis. Adm. Code. Ongoing maintenance is needed to contain these populations so they do not re-establish throughout the waterbody, spread to other waters, or impair navigation and other beneficial uses of the waterbody.

Funding Possibilities

Maximum amount of grant funding will be determined by DNR based on the sponsor's permit application fee, specified monitoring and reporting requirements in the permit, or DNR-approved management plan. The maximum grant amount shall not exceed the cost of the permit application fee.

LAKE MANAGEMENT PLANNING GRANTS

Lake management planning grants are intended to provide financial assistance to eligible applicants for the collection, analysis, and communication of information needed to conduct studies and develop management plans to protect and restore lakes and their watersheds. Projects funded under this subprogram often become the basis for implementation projects funded with Lake Protection grants. There are two categories of lake management planning grants: small-scale and large-scale.

SMALL SCALE LAKE MANAGEMENT PROJECTS

Small-scale projects are intended to address the planning needs of lakes where education, enhancing lake organizational capacity, and obtaining information on specific lake conditions are the primary project objectives. These grants are well suited for beginning the planning process, conducting minor plan updates, or developing plans and specification for implementing a management recommendation. Eligible projects include:

- Collect and report chemical, biological, and physical data about lake ecosystems for a Tier I assessments, Tier II diagnostic or Tier III project evaluation.
 - Tier I if initial basic monitoring is needed to assess the general condition or health of the lake.
 - Tier II if an assessment has been conducted and more detailed data collection is needed to diagnose suspected problems and identify management options.
 - Tier III if the monitoring and assessment will be used to evaluate the effectiveness of a recently implemented project or lake management strategy.
- Collecting and disseminating existing information about lakes for the purpose of broadening the understanding of lake use, Lake Ecosystem conditions and lake management techniques.
- Conducting workshops or trainings needed to support planning or project implementation.
- Projects that will assist management units as defined in s. NR191.03 (4) & s. NR 190.003 (4) the formation of goals and objectives for the management of a lake or lakes.

Funding Possibilities

Maximum amount of grant funding is 67% of the total project costs, not to exceed \$3,000.

LARGE SCALE LAKE MANAGEMENT PROJECTS

Large-scale projects are intended to address the needs of larger lakes and lakes with complex and technical planning challenges. The result will be a lake management plan; more than one grant may be needed to complete the plan. Eligible projects include:

- Collection of new or updated, physical, chemical and biological information about lakes or lake ecosystems.
- Definition and mapping of Lake Watershed boundaries, sub-boundaries and drainage system components.
- Descriptions and mapping of existing and potential land conditions, activities and uses within lake watersheds that may affect the water quality of a lake or its ecosystem.
- Assessments of water quality and of fish, aquatic life, and their habitat.
- Institutional assessment of lake protection regulations - review, evaluation or development of ordinances and other local regulations related to the control of pollution sources, recreational use or other human activities that may impact water quality, fish and wildlife habitat, natural beauty or other components of the lake ecosystem.
- Collection of sociological information through surveys or questionnaires to assess attitudes and needs and identify problems necessary to the development of a long-term lake management plan.
- Analysis, evaluation, reporting and dissemination of information obtained as part of the planning project and the development of management plans.
- Development of alternative management strategies, plans and specific project designs, engineering or construction plans and specifications necessary to identify and implement an appropriate lake protection or improvement project.

Funding Possibilities

Maximum amount of grant funding is 67% of the total project costs, not to exceed \$25,000. Multiple grants in sequence may be used to complete a planning project, not to exceed \$100,000 for each lake. The maximum grant award in any one year is \$50,000 for each lake. If phasing is necessary, all phases should be fully identified and a timeline identified in the initial application.

LAKE PROTECTION GRANTS

Lake protection and classification grants assist eligible applicants with implementation of lake protection and restoration projects that protect or improve water quality, habitat or the elements of lake ecosystems. There are four basic Lake Protection subprograms: a) Fee simple or Easement Land Acquisition b) Wetland and Shoreline Habitat Restoration c) Lake Management Plan Implementation d) Healthy Lakes Projects.

HEALTHY LAKES PROJECTS

The Healthy Lakes grants are a sub-set of Plan Implementation Grants intended as a way to fund increased installation of select best management practices (BMPs) on waterfront properties without the burden of developing a complex lake management plan. Details on the select best practices can be found in the Wisconsin Healthy Lakes Implementation Plan and best practice fact sheets.

Eligible best practices with pre-set funding limits are defined in the Wisconsin Healthy Lakes Implementation Plan, which local sponsors can adopt by resolution and/or integrate into their own local planning efforts. By adopting the Wisconsin Healthy Lakes Implementation Plan, your lake organization is immediately eligible to implement the specified best practices. Additional technical information for each of the eligible practices is described in associated factsheets. The intent of the Healthy Lakes grants is to fund shovel-ready projects that are relatively inexpensive and straight-forward. The Healthy Lakes grant category is not intended for large, complex projects, particularly those that may require engineering design. All Healthy Lake grants have a standard 2-year timeline.

Funding Possibilities

Maximum amount of grant funding is 75% of the total project cost, not to exceed \$25,000. Grants run for a 2-year time period. Maximum costs per practice are also identified in the Wisconsin Healthy Lakes Implementation Plan.

Works Cited (Not final yet)

- Christensen, D. H. (1996). Impacts of lakeshore residential development on coarse woody debris in north temperate lakes. *Ecological Applications* 6 (4), 1143-1149.
- Eichler, L. B. (1993). Suction harvesting of EWM and its effect on native plant communities. *Journal of Aquatic Plant Management* 31, 144-148.
- Gettsinger, K. T. (1997). Restoring native vegetation in a Eurasian water milfoil dominated plant community using the herbicide triclopyr. *Regulated Rivers: Research and Management* 13, 357-375.
- Glomski, L. M. (2010). Response of Eurasian and hybrid watermilfoil to low use rates and extended exposures of 2, 4-D and triclopyr. *Journal of Aquatic Plant Management* 48 (12).
- Jennings, M. E. (2003). Is littoral habitat affected by residential development and land use in watersheds of Wisconsin lakes? *Lake Reservoir Management*, 19 (3), 272-279.
- Kelting, D. a. (2010). Cost and effectiveness of hand harvesting to control the EWM population in Upper Saranac Lake, New York. *Journal of Aquatic Plant Management* 48.
- LaRue, E. Z. (2012). *Hard to Kill: Hybrid Watermilfoil are less sensitive to a commonly used herbicide.*
- Madsen, J. (1997). *Methods for management of nonindigenous aquatic plants.* New York: Springer.
- Madsen, J. (2000). *Advantages and disadvantages of aquatic plant management techniques.* Vicksburg, MS: US Army Corps of Engineers Aquatic Plant Control Research Program.
- Newman, R. H. (1996). Effects of the potential biological control agent, *Euhrychiopsis lecontei*, on EWM in experimental tanks. *Aquatic Botany* 53, 131-150.
- Parepa, M., Schaffner, U., & Bossdorf, O. (2012). Sources and modes of action of invasive knotweed allelopathy: the effects of leaf litter and trained soil on the germination and growth of native plants.
- Pine, R. a. (1991). Plant preferences of Triploid grass carp. *Journal of Aquatic Plant Management* 29, 80-82.
- Poovey, A. S. (2007). Susceptibility of EWM (*Myriophyllum spicatum*) and a milfoil hybrid (*M. spicatum* x *M. sibiricum*) to triclopyr and 2, 4-D amine. *Journal of Aquatic Plant Management* 45, 111-115.
- Sorsa, K. N. (1988). Integrated control of Eurasian water milfoil by a fungal pathogen and herbicide. *Journal of Aquatic Plant Management* 26, 12-17.
- Tobiessen, P. S. (1992). Dredging to control curly-leaf pondweed: a decade later. *Journal of Aquatic Plant Management* 30, 71-72.
- Wolter, M. (2012). *Lakeshore Woody Habitat in Review.* Hayward, WI: Wisconsin Department of Natural Resources.

Appendix B
NR 107

Appendix C
WDNR Dredging Checklist

Appendix D
WDNR Lake Shoreland and Shallows Habitat Monitoring Field Protocol

Appendix E
Round Lake Aquatic Plant Management Goals, Objectives, and Actions

Appendix F
Funding and Implementation Matrix

Appendix G
Calendar of Actions

